

Decision Theoretic Generalizations of the PAC Model for Neural Net and Other Learning Applications

DAVID HAUSSLER

*Baskin Center for Computer Engineering and Information Sciences,
University of California, Santa Cruz, California 95064*

We describe a generalization of the PAC learning model that is based on statistical decision theory. In this model the learner receives randomly drawn examples, each example consisting of an instance $x \in X$ and an outcome $y \in Y$, and tries to find a decision rule $h: X \rightarrow A$, where $h \in \mathcal{H}$, that specifies the appropriate action $a \in A$ to take for each instance x in order to minimize the expectation of a loss $l(y, a)$. Here X , Y , and A are arbitrary sets, l is a real-valued function, and examples are generated according to an arbitrary joint distribution on $X \times Y$. Special cases include the problem of learning a function from X into Y , the problem of learning the conditional probability distribution on Y given X (regression), and the problem of learning a distribution on X (density estimation). We give theorems on the uniform convergence of empirical loss estimates to true expected loss rates for certain decision rule spaces \mathcal{H} , and show how this implies learnability with bounded sample size, disregarding computational complexity. As an application, we give distribution-independent upper bounds on the sample size needed for learning with feedforward neural networks. Our theorems use a generalized notion of VC dimension that applies to classes of real-valued functions, adapted from Vapnik and Pollard's work, and a notion of *capacity* and *metric dimension* for classes of functions that map into a bounded metric space. © 1992 Academic Press, Inc.

1. INTRODUCTION

The introduction of the Probably Approximately Correct (PAC) model (Valiant, 1984; Angluin, 1988) of learning from examples has done an admirable job of drawing together practitioners of machine learning with theoretically oriented computer scientists in the pursuit of a solid and useful mathematical foundation for applied machine learning work. These practitioners include both those in mainstream artificial intelligence and those in neural net research. However, in attempts to address the issues that are relevant to this applied work in machine learning, a number of shortcomings of the model have cropped up repeatedly. Among these are the following:

1. The model is defined only for $\{0, 1\}$ -valued functions. Practitioners would like to learn functions on an instance space X that take

values in an arbitrary set Y ; e.g., multi-valued discrete functions, real-valued functions, and vector-valued functions.

2. Some practitioners are wary of the assumption that the examples are generated from an underlying “target function,” and are not satisfied with the noise models that have been proposed to weaken this assumption (e.g., Angluin and Laird, 1988; Sloan, 1988; Shackelford and Volper, 1988). They would like to see more general regression models investigated in which the y component in a training example $(x, y) \in X \times Y$ is randomly specified according to a conditional distribution on Y , given x . Here the general goal is to approximate this conditional distribution for each instance $x \in X$. In the computational learning theory literature, a model of this type is investigated in Kearns and Schapire (1990), with $Y = \{0, 1\}$, and for a more general case in Yamanishi (1990).

3. Many learning problems are unsupervised; i.e., the learner has access only to randomly drawn, unlabeled examples from an instance space X . Here learning can often be viewed as some form of approximation of the distribution that is generating these examples. This is usually called *density estimation* when the instance space X is continuous and no specific parametric form for the underlying distribution on X is assumed. It is often called *parameter estimation* when specific parametric probability models are used. One example of this in the computational learning theory literature is the recent investigation of Abe and Warmuth (1990) into the complexity of learning the parameters in a hidden Markov model.

Our purpose here is twofold. First, we propose an extension of the PAC model, based on the work of Vapnik and Chervonenkis (Vapnik, 1989) and Pollard (1984, 1990), that addresses these and other issues. Second, we use this extension to obtain distribution-independent upper bounds on the size of the training set needed for learning with various kinds of feed-forward neural networks (Rumelhart and McClelland, 1986; Poggio and Girosi, 1989), a popular learning method that is not covered by the basic PAC model.

1.1. Overview of the Proposed Framework

To extend the PAC model, we propose a more general framework based on statistical decision theory (see, e.g., Ferguson (1967), Kiefer (1987), or Berger (1985)). In this general framework we assume the learner receives randomly drawn training examples, each example consisting of an instance $x \in X$ and an outcome $y \in Y$, where X and Y are arbitrary sets called *instance* and *outcome spaces*, respectively. These examples are generated according to a joint distribution on $X \times Y$, unknown to the learner. This distribution comes from a (known) class \mathcal{P} of joint distributions on $X \times Y$,

representing possible “states of nature.” After training, the learner receives further random examples drawn from this same joint distribution. For each example (x, y) , the learner will be shown only the instance x . Then he will be asked to choose an action a from a set of possible actions A , called the *decision space*. Following this, the outcome y will be revealed to the learner. In the case that we examine here, the outcome y depends only on the instance x and not on the action a chosen by the learner. For each action a and outcome y , the learner will suffer a loss, which is measured by a fixed real-valued *loss function* l on $Y \times A$. We assume that the loss function is known to the learner. The learner tries to choose his actions so as to minimize his loss.

Here we look at the case in which, based on the training examples, the learner develops a deterministic strategy that specifies what he believes is the appropriate action a for each instance x in X . He then uses this strategy on all future examples. Thus we look at “batch” learning rather than “incremental” or “on-line” learning (Littlestone, 1988). The learner’s strategy, which is a function from the instance space X into the decision space A , is called a *decision rule*. We assume that the decision rule is chosen from a fixed *decision rule space* \mathcal{H} of functions from X into A . For example, instances in X may be encoded as inputs to a neural network, and outputs of the network may be interpreted as actions in A . In this case the network represents a decision rule, and the decision rule space \mathcal{H} may be all functions represented by networks obtained by varying the parameters of a fixed underlying network. The goal of learning is to find a decision rule in \mathcal{H} that minimizes the expected loss, when examples are drawn at random from the unknown joint distribution on $X \times Y$.

This learning framework can be applied in a variety of situations. We now give several illustrations. For further discussion, we refer the reader to the excellent surveys of White (1990b), Barron (1989), Devroye (1988), and Vapnik (1989), to which we are greatly indebted. We also recommend the text by Kiefer (1987), for a general introduction to statistical inference and decision theory.

1.1.1. *Betting Example*

For our first example, consider the problem of learning to maximize profit (or minimize loss!) at the horse races. Here an instance x in X is a race, an action a in A consists of placing or not placing a certain bet, and an outcome y in Y is determined by the winner and the second and third place finishers. The loss $l(y, a)$ is the amount of money lost when bet a is placed and the outcome of the race is y . A negative loss is interpreted as a gain. The joint distribution on $X \times Y$ represents the probability of various races and outcomes. This joint distribution is unknown to the learner; he only has random examples $(x_1, y_1), \dots, (x_m, y_m)$, each consisting

of a race/outcome pair generated from this distribution. From these examples, the learner develops a deterministic betting strategy (decision rule). The best decision rule h is one that specifies a bet a for each race x that minimizes the expectation of the loss $l(y, a)$, when y is chosen randomly from the unknown conditional distribution on Y given x , which is determined by the underlying joint distribution on $X \times Y$. This (not necessarily unique) best decision rule minimizes the expected loss on a random example (x, y) . It is known as the *Bayes optimal decision rule*. The learner tries to approximate the Bayes optimal decision rule as best he can using decision rules from a given decision rule space \mathcal{H} (e.g., “simple” or “easy to compute” decision rules, or perhaps decision rules that can be represented by a particular kind of neural network).

1.1.2. Classification

As a second example, consider the problem of medical diagnosis. Here an instance x is a vector of measurements from medical tests conducted on the patient, an action a is a diagnosis of the patient’s disease state, and an outcome y may be defined as the actual disease state of the patient. Here $A = Y$; i.e., the possible diagnoses are the same as the possible disease states. To specify the loss function l , we may stipulate that there is zero loss for the correct diagnosis $a = y$, but for each pair (y, a) with diagnosis a differing from disease state y there is some positive real loss $l(y, a)$, depending on the severity of the consequences of that particular misdiagnosis. Here a decision rule is a diagnostic method, and Bayes optimal decision rule is the one that minimizes the expected loss from misdiagnosis when examples (x, y) of test results and associated disease states occur randomly according to some unknown “natural” joint distribution.

This medical diagnosis situation is a typical example of a *classification learning* problem in the field of pattern recognition (see, e.g., Duda and Hart (1973)). The problem of learning a Boolean function from noise-free examples, as investigated in the PAC model, is a special case of classification learning. Here the outcome space Y is $\{0, 1\}$ and only the instance x in an example (x, y) is drawn at random. The outcome y is $f(x)$ for some unknown Boolean *target function* f , rather than being determined stochastically. As above, the decision space A is the same as the outcome space Y , and the action a can be interpreted as a prediction of the outcome y . Hence, a decision rule h maps from the instance space X into the outcome space Y , just as the target function does. In much of AI, and in PAC learning in particular, it is common to refer to h as a *hypothesis* in this case, and to \mathcal{H} as the *hypothesis space*.

This same setup, where the outcome y is a function of the instance x , can be applied to any function learning problem by letting X and Y be

arbitrary sets. In the general function learning problem, the loss function $l(y, a)$ usually measures the distance between the prediction a and the actual value y in some metric. In the PAC model, l is the discrete metric: $l(y, a) = 0$ if $a = y$, else $l(y, a) = 1$. Thus the expected loss of the decision rule (or hypothesis) is just the probability that it predicts incorrectly, the usual PAC notion of the *error* of the hypothesis. In general, Y may be a set of strings, graphs, real vectors, etc., in which case other distance metrics or more general kinds of loss functions may be more appropriate.

1.1.3. Regression

The general problem of regression has a different character from that of classification learning, but can also be addressed in the decision theoretic learning framework. To illustrate this, as a third example consider a variant of the medical diagnosis situation in which the doctor provides an estimate of the probability that the patient has each of several diseases, rather than predicting that he has one specific disease or asserting that he is healthy. (Here we assume that the actual disease state includes at most one disease.) For example, the doctor may say "Given these test results x , I would say you have disease 1 with probability 55%, disease 2 with probability 5%, and no disease at all with probability 40%." Here the doctor is actually trying to estimate the conditional distribution on disease states Y given the test results x . Her action a entails providing a vector of parameters that determine that estimated distribution, e.g., (0.55, 0.05, 0.4). The decision space A is the set of all such parameter vectors.

Now let Y be an arbitrary discrete outcome space. Keeping the instance x fixed, for each parameter vector a in A and outcome y in Y let $\hat{P}(y; a)$ denote the probability of outcome y with respect to the distribution on Y defined by the parameter vector a . Thus when we take action a on instance x , we are asserting that, given the instance x , we estimate the conditional probability of outcome y to be $\hat{P}(y; a)$ for each outcome y in Y . Let $P(y)$ denote the actual conditional probability of outcome y , given the instance x , with respect to the unknown joint distribution on $X \times Y$. (The distributions P and \hat{P} can be replaced by densities when Y is continuous.) Let us define¹ the loss function l by setting $l(y, a) = -\log \hat{P}(y; a)$. This is called the (negative) *log likelihood* loss function. If we define loss in this way, then the expected loss resulting from action a has a natural information theoretic interpretation:² it is the *Kullback–Leibler divergence* (Kullback,

¹ We assume $\hat{P}(y; a) > 0$ for all y in Y .

² The Kullback–Leibler divergence from P to \hat{P} , denoted $I(P \| \hat{P})$, is defined as $\sum_{y \in Y} P(y) \log(P(y)/\hat{P}(y; a))$ for countable Y . The entropy of P , denoted $H(P)$, is $-\sum_{y \in Y} P(y) \log P(y)$. Thus $I(P \| \hat{P}) + H(P) = -\sum_{y \in Y} P(y) \log \hat{P}(y; a)$, which is the expectation of the (negative) log likelihood loss. Analogous results hold for densities when the relevant quantities are finite (Kullback, 1959).

1959) (or *information gain* (Renyi, 1970)) from the actual conditional probability distribution P to the estimated conditional distribution \hat{P} , plus the entropy of P .

For a given x , the entropy of the true conditional distribution P is a constant, independent of the action a . Thus choosing the action a for each instance x that minimizes the expected log likelihood loss is equivalent to choosing the action a that gives the closest estimate \hat{P} to the true conditional distribution P over possible outcomes in Y as measured by the Kullback–Leibler divergence, given that instance x . It is well known that the Kullback–Leibler divergence is minimized when $\hat{P} = P$. This is the Bayes optimal decision rule in regression.

In the regression version of our medical diagnosis situation, the definition of the log likelihood loss function depends on the interpretation of the components of the parameter vector a . If there are k possible diseases and the patient can have at most one of these, then we might have $k + 1$ possible mutually exclusive disease states y_1, \dots, y_{k+1} , where y_{k+1} means healthy. Hence $Y = \{y_1, \dots, y_{k+1}\}$. Then we might specify that an action a takes the form

$$a = (a_1, \dots, a_{k+1}),$$

where $a_i = \hat{P}(y_i; a)$, the estimated probability of disease state y_i . Here the components of the vector a must be positive and sum to one. In this case the log likelihood loss would be $l(y_i, a) = -\log a_i = -\log \hat{P}(y_i; a)$.

Often the constraints on the components of a are a nuisance, so other interpretations of a are used; e.g., that $a_i = \log \hat{P}(y_i; a) - \log \hat{P}(y_{k+1}; a)$ for each i , $1 \leq i \leq k + 1$. In this case the a_1, \dots, a_k are arbitrary real numbers and $a_{k+1} = 0$, and hence can be ignored. Since $\hat{P}(y_i; a) = e^{a_i} / \sum_{j=1}^{k+1} e^{a_j}$, the log likelihood loss is $l(y_i, a) = -a_i + \log \sum_{j=1}^{k+1} e^{a_j} = -a_i + \log(1 + \sum_{j=1}^k e^{a_j})$. This is known as the *logistic loss* (McCullagh and Nelder, 1989; Barron, 1989).

A third interpretation would be to allow the possibility that the patient may have more than one disease, and assume, for the purposes of estimation, that diseases occur independently. Then the disease state y might be defined as a binary vector of length k , where the i th bit y_i is 1 if and only if the i th disease is present. Hence $Y = \{0, 1\}^k$. Similarly, the vector a would be a vector of independent probabilities (a_1, \dots, a_k) , where a_i is the estimated probability of the patient having the i th disease. In this case

$$\hat{P}(y; a) = \prod_{i=1}^k a_i^{y_i} (1 - a_i)^{(1 - y_i)}$$

and the log likelihood loss is

$$l(y, a) = - \sum_{i=1}^k (y_i \log a_i + (1 - y_i) \log(1 - a_i)),$$

which we call the *cross entropy loss*.

In the medical diagnosis example, the outcome space Y is discrete. However, in most uses of regression Y is real valued; i.e., the outcome y is the measurement of some real valued quantity, and the instance x represents the experimental conditions under which this quantity was measured. In this case regression is usually defined as estimating the conditional expectation of Y given the instance x . Thus $A \subset \mathfrak{R}$, and the action $a \in A$ for a given instance x consists of an estimate of the mean of the various outcomes y that would typically be observed for that instance x . It is easy to show that by using the *quadratic* loss function $l(y, a) = (a - y)^2$, the expected loss is minimized when a is the true mean, and hence this version of regression also fits naturally³ into the decision theoretic framework. An alternate approach is to use the L_1 loss function $l(y, a) = |a - y|$, in which case the expected loss is minimized when a is the median of the conditional distribution of Y given the instance x . (See, e.g., White, 1990b; Haussler, 1990.)

1.1.4. Density and Parameter Estimation

Finally, the problems of parameter estimation and density estimation can also be viewed as special cases of this decision theoretic framework. For parameter estimation, note that when the instance space X has only one element, the particular instance x can be ignored entirely. Thus the regression problem reduces to the problem of estimating the parameters of a single distribution on the outcome space Y from a sample of random outcomes y from Y , i.e., to the simpler problem of parameter estimation. Here

³ In fact, the standard version of regression, defined as estimating the conditional mean of Y given instance x using the quadratic loss function, is actually a special case of the general version of regression defined above, where for continuous outcome spaces Y , the object is to estimate the parameters specifying the conditional density of Y given instance x , using the log likelihood loss function. To see this, assume that we represent the conditional density on Y with a Gaussian density $\hat{p}(y; \mu, \sigma) = (2\pi\sigma^2)^{-1/2} e^{-(\mu - y)^2/2\sigma^2}$, where μ is the mean and σ^2 the variance. Let the variance be fixed, independent of x , so that the estimate $\hat{p}(y; \mu, \sigma)$ of the conditional density on Y given x is completely determined by the mean μ . Thus the decision space $A \subset \mathfrak{R}$, and each action a in A is interpreted as specifying the mean of a Gaussian density. Substituting $\mu = a$ and evaluating $-\log \hat{p}(y; \mu, \sigma)$, the log likelihood loss is seen to be $l(y, a) = (1/2\sigma^2)(a - y)^2 + \frac{1}{2} \log(2\pi\sigma^2)$. For fixed variance σ^2 , this is equivalent, for learning, to the quadratic loss $(a - y)^2$, since additive and multiplicative constants in the definition of l only rescale it without changing the value of a that minimizes its expectation.

the decision rule is not a function but merely a single vector of parameters, and the decision rule space \mathcal{H} is the same as the decision space A .

For density estimation, we can consider the dual case in which the outcome space Y has only one element, and hence can be ignored. Thus examples are unlabeled instances x drawn randomly from some density $p(x)$ on X . Let the decision set A be the positive real numbers and let each decision rule h in \mathcal{H} be a density on X . Then, as above, information theoretic considerations suggest the loss function $l(y, a) = l(a) = -\log a$. Again, as above, the expected loss of h is minimized when h is the true density p . Further, if p is not a member of \mathcal{H} , then the best decision rule in \mathcal{H} , in terms of minimizing the expected loss, is the one with the smallest Kullback–Leibler divergence from the true density p (Kullback, 1959).

When the instance space X is discrete, we are not estimating a density on X but rather a probability distribution. The same ideas as above carry over, except that we let the decision space $A = (0, 1)$ and each decision rule h in \mathcal{H} represents a probability distribution on X . Here we can also use the same loss function, and it has the same properties.

These examples illustrate the diversity of the learning problems that can be cast in the proposed decision theoretic framework, even under the restrictive assumptions we make here, i.e., that the outcome y does not depend on the action a , and that the learner always observes both the outcome and the loss. By weakening these assumptions, we can model other types of learning as well, including *associative reinforcement learning* (Barto and Anandan, 1985; Gullapalli, 1990) and the theory of *learning automata* (with static environment) (Narendra and Thathachar, 1989). However, we will not pursue this here.

1.2. Summary and Discussion of the Results Presented Here

There are three major practical issues in this decision theoretic view of learning. The first is the number of random examples needed in order to be able to produce a good decision rule in the decision rule space \mathcal{H} , i.e., a decision rule whose expected loss is near the minimum of all decision rules in \mathcal{H} . If too few examples are used, we run into the problem of *overfitting*, where the decision rule produced performs well on the training data, but not on further random examples drawn from the same joint distribution that generated this training data. The second is the adequacy of the decision rule space \mathcal{H} . If \mathcal{H} does not contain any decision rule with expected loss close to that of Bayes optimal decision rule for the particular joint distribution we are dealing with, then we can never hope to achieve near optimal performance using this decision rule space. Choosing the right decision rule space often requires considerable insight into the particular problem domain. Finally, the third practical problem is the computational

complexity of the method we use to produce our decision rule from the training examples. This issue has been addressed extensively in the PAC literature, and is also addressed in Kearns and Schapire (1990), and Abe and Warmuth (1990). Of these three important issues, here we examine only the first. This issue is referred to as the problem of estimating the “sample complexity” of the learning problem in the PAC literature (Ehrenfeucht *et al.*, 1989).

The number of random training examples needed to avoid overfitting depends critically on the nature of the decision rule space used. Different kinds of decision rule spaces are used in different areas of learning research, partly because different kinds of instance and outcome spaces are used. In pattern recognition and statistics, the instance space X is usually a finite dimensional real vector space; i.e., each instance consists of a vector of real valued measurements of some attributes. In density estimation, a decision rule represents a density on X , and many choices are possible. One common choice is a mixture of Gaussian densities (e.g., Duda and Hart, 1973; Nowlan, 1990). In standard regression, the outcome and decision spaces Y and A are identical and real valued, and linear functions are most often used as decision rules. For more complex outcome spaces, such as those in the medical diagnosis example given above, the decision rule space for regression is usually defined using a *generalized linear model* (McCullagh and Nelder, 1989). Similarly, in binary classification, where there are only two possible outcomes in Y as in the PAC model, linear threshold functions are most often used as decision rules, and there are straightforward generalizations for the case of k -ary classification (see, e.g., Duda and Hart, 1973). This “linear bias” in pattern recognition and statistics is in contrast to that in the PAC model and other AI areas, including work in neural networks, in which a rich variety of decision rule spaces are used (see, e.g., Touretsky, 1989, 1990; Haussler, 1988, 1989). Our main goal here is to develop analytic tools to help understand the problem of overfitting in these more complex decision rule spaces.

In order to focus on the problem of overfitting, we take a simplified view of learning, in which the learner chooses a decision rule space \mathcal{H} , and then tries to find a decision rule in \mathcal{H} with near minimal expected loss. To do this, the learner looks for a decision rule that minimizes the observed average loss on the training examples, which is called *empirical loss* or *empirical risk*. For example, in standard linear regression⁴ the learning algorithm is the method of least squares; i.e., we find the linear function h that minimizes the average of $l(y, h(x)) = (h(x) - y)^2$ over all examples

⁴ For general regression with the negative log likelihood loss function, the principle of minimizing empirical loss is the same as the principle of *maximum likelihood* (Berger, 1985; Kiefer, 1987).

(x, y) in our training set. It is well known that if we have too few training examples, then we tend to overfit them, and the function we find does not come close to minimizing the actual expected quadratic loss, which would be obtained by integrating over all possible (mostly unseen) examples with respect to the unknown joint distribution on them. This same situation occurs with all nontrivial decision rule spaces, including the nonlinear regression models defined by feedforward neural nets.

Using certain measures of the “dimension” or “capacity” of the decision rule space \mathcal{H} and classes derived from \mathcal{H} (see below), we obtain general upper bounds on the number of random training examples needed so that with high probability, any decision rule in \mathcal{H} that has small empirical loss on the training examples will have small actual expected loss; i.e. we get uniform convergence results for empirical estimates like those in Vapnik (1982), Dudley (1984), and Pollard (1984, 1990). We show how these give upper bounds on sufficient training sample size like those derived in Blumer *et al.* (1989) and elsewhere using the notion of the VC dimension, and generalize those results.

As an application, we give specific bounds on the number of training examples needed to avoid overfitting when learning with the decision rule space of feedforward neural nets (Rumelhart and McClelland, 1986), extending previous work in Baum and Haussler (1989) and White (1990a) (see also related work in Anthony and Shawe-Taylor, 1990). These are the nets most widely used in current neural net learning research. Our model for feedforward neural nets is quite general in that it allows many types of units in the nets, including quasi-linear units (Rumelhart and McClelland, 1986), radial basis units (Poggio and Girosi, 1989), and product units (Durbin and Rumelhart, 1989).

In our general setting, successful learning means finding a decision rule with average loss close to minimal over all decision rules in the given decision rule space, rather than loss close to zero as in the PAC model. In addition to using an additive model as in Lineal *et al.* (1988), we also define “close to” using a measure of relative difference (the d , *metric*) similar to the standard multiplicative measure of approximation used in combinatorial optimization. This allows us to state the relevant uniform convergence bounds as generalized “Chernoff-style” (Angluin and Valiant, 1979) bounds, as in Pollard (1986) and Breiman *et al.* (1984, Chap. 12), rather than “Hoeffding-style” bounds (as in Pollard’s (1984) results) giving better bounds on sufficient training sample size in some important cases. These two types of bounds are analogous to the two types of bounds that Vapnik (1982) gives in his book, in that one uses a measure of absolute difference and the other a measure of relative difference. However, both of our bounds are “two-sided”; i.e., they bound deviations both above and below the mean.

We give these upper bounds on required sample size only to give some indication of the order-of-magnitude dependence of sample size on certain critical parameters of the learning problem, and to illustrate the theory. They are still too crude to be used directly in practice, e.g., as explicit formulae for choosing an appropriate sample size. Cross validation techniques, in which some of the training examples are held in reserve and used instead to test the performance of the decision rules produced by the learning algorithm, are likely to perform better for this task in practice (see, e.g., White, 1990a; Weiss and Kulikowski, 1991). Nevertheless, cross validation is only a means of estimating the amount of overfitting in the learning method in particular cases; i.e. it is only an engineering trick and provides no scientific explanation of the phenomenon. Our goal is to understand and explain overfitting in general decision rule spaces, from a scientific rather than an engineering viewpoint.

Finally, we should note that in practice, many learning algorithms do more than just search for a decision rule in a fixed decision rule space that minimizes empirical loss. For example, it is common to let the decision rule space depend on the number of training examples available, using richer and richer decision rule spaces as more examples become available (see, e.g., White, 1990a; Blumer *et al.*, 1989). This can allow the learning algorithm to produce a sequence of decision rules with expected losses that approach the loss of the Bayes optimal decision rule in the limit of infinite training sample size for a large class of possible joint distributions. The results given here can be used to estimate the appropriate rate at which the decision rule space should grow relative to the sample size to avoid overfitting. Other approaches, e.g., the method of *structural risk minimization* introduced by Vapnik (1982), and the *Bayesian* (Berger, 1985; Mackay, 1992; Buntine and Weigend, 1991) and *minimum description length* (MDL) approaches (Barron and Cover, 1990; Rissanen, 1986), try to find a decision rule that minimizes some function of empirical loss and decision rule complexity. These can also achieve expected loss approaching that of Bayes optimal decision rule in the limit, and may be more effective in practice. Although uniform convergence results such as those we develop here are also used in the analysis of such methods (Vapnik, 1982) (and in the analysis of cross-validation methods (Nolan and Pollard, 1987)), the full treatment of such approaches is beyond the scope of the present paper. It should also be noted that Bayesian methods and structural risk minimization can be applied even when the decision rule space includes only neural networks of a fixed size. An example is the recent work using weight penalty functions in neural net training (Weigend *et al.*, 1990; LeCun *et al.*, 1990; Nowlan and Hinton, 1991; MacKay, 1992; Buntine and Weigend, 1991). Such approaches may significantly reduce the training sample size needed to avoid overfitting in practice.

1.3. Overview of Methods Used

We now briefly discuss the methodology and previous work used in obtaining our results. Our work builds directly on the work of Vapnik and Chervonenkis, Pollard, and Dudley on the uniform convergence of empirical estimates (Vapnik, 1982; Pollard, 1984; Dudley, 1984) and its application to pattern recognition (Vapnik, 1982, 1989; Devroye, 1988). It also builds on the work of Benedek and Itai (1988) on PAC learnability with respect to specific probability distributions, and is related to the work of Natarajan and Tadeipalli on extensions of the VC dimension to multivalued functions (Natarajan and Tadeipalli, 1988; Natarajan, 1989b) and PAC learnability with respect to classes of probability distributions (Natarajan, 1988, 1989a). In addition, Quiroz (1989) and Kulkarni (1989) have each independently generalized the PAC model in a related manner.

One of the key ideas we use is the notion of an ε -cover of a metric space (Dudley, 1989; Pollard, 1984; Benedek and Itai, 1988; Natarajan, 1989a, Quiroz, 1989) and the associated idea of *metric dimension* (Kolmogorov and Tihomirov, 1961) (also called the *fractal dimension* (Farmer, 1982)). This notion of dimension has played an important role in the now very active study of fractals in nature (Mandelbrot, 1982), especially in connection with chaos in dynamical systems (Farmer, 1982; Farmer *et al.*, 1983). Here we build further on the beautiful results of Vapnik and Chervonenkis (Vapnik, 1982), Dudley (1978), and Pollard (1984), which relate a type of generalized VC dimension for a decision rule space to the number of balls of radius ε required to cover the space, with respect to certain metrics. The sizes of the smallest such covers determine the *metric dimension* of the space. Our treatment closely parallels the approach given in Pollard (1990). It is interesting to note that related results connecting ε -covers with the VC dimension have also been independently developed in Benedek and Itai (1988) and in recent computational geometry work (Welzl, 1988).⁵ This work seems to lead to a potentially rich area of investigation that combines elements of combinatorics, topology and geometry, and probability and measure in a novel framework. We feel that this area is not only fascinating from a purely mathematical standpoint, but also potentially very useful in machine learning and other applied fields.

1.4. Organization of the Paper

The remainder of the paper is organized as follows. The learning framework we described above in Section 1.1 is defined more formally in

⁵ Specifically, Lemma 7.13 of Dudley (1978) is nearly equivalent to Lemma 4.1 of Welzl (1988) (using the primal space instead of the dual). This result also gives a stronger version of Theorem 4, part (3) of Benedek and Itai (1988). We give a still stronger version of this result in Theorem 6 below.

Section 2. There we also look at the question of evaluating the performance of learning algorithms in terms of the number of training examples they use. This question is also formalized from a decision theory perspective. We then provide a lemma (Lemma 1) that can be used to evaluate the performance of learning algorithms that work by minimizing empirical loss. To use this lemma, we need bounds on the rate of uniform convergence of empirical loss estimates to true expected losses. These are given in Section 3. The key bound is given in Theorem 2 in Section 3, and in a more general version in Theorem 3.

To use the bound from Theorem 2 we need bounds on the “random covering numbers” associated with the decision rule space \mathcal{H} , the loss function l , and the distribution P . These are related to the idea of an ε -cover described above. In Section 4 we introduce Pollard’s notion of the pseudo dimension as a means of bounding the random covering numbers. Applications of this method to several learning problems are described in Section 5.

The techniques of Sections 4 and 5 only apply to the case when the action set A is real-valued. Tools for bounding the random covering numbers that apply in more general cases are developed in Section 6. Here we introduce the notion of the capacity of the decision rule space \mathcal{H} (for a particular loss function l), and the related notion of the metric dimension of \mathcal{H} . In Section 7 we use these notions to obtain bounds on the performance (in terms of the number of training examples used) of learning algorithms that use multilayer feedforward neural networks, and work by minimizing empirical loss (Corollary 3). Finally, some further discussion of our results is given in the conclusion, Section 8.

Many of the more technical proofs and definitions have been moved into the Appendix to make the paper more readable. The Appendix has several sections. Section 9.1 contains a brief overview of the theory of metric spaces, ε -covers, and metric dimension. Notation from this section is used in several places in the paper. Section 9.2 deals with certain technical measurability requirements. Section 9.3 gives an analogue of Chernoff and Hoeffding bounds using the d_v metric. Section 9.4 contains the proof of Theorem 2. Finally, Section 9.5 contains a result on feedforward neural networks of linear threshold functions that is similar to that given in Baum and Haussler (1989), and provides a counterpart to Corollary 3 in Section 7.

1.5. Notational Conventions

We denote the real numbers by \mathfrak{R} and the non-negative real numbers by \mathfrak{R}^+ . By \log and \ln we denote the logarithm base 2 and the natural

logarithm, respectively. We use $E(\cdot)$ to denote the expectation of a random variable, and $\text{Var}(\cdot)$ to denote the variance of a random variable. When the probability space is defined implicitly from the context, we use $\text{Pr}(\cdot)$ to denote the probability of a set. However, usually the measure on the underlying probability space is defined explicitly using the symbol P .

Here, P usually denotes a probability measure on some appropriate⁶ σ -algebra over the set $Z = X \times Y$, where X is the instance space and Y is the outcome space. We use P^m to denote the m -fold product measure on Z^m . Functions on Z and subsets of Z mentioned in what follows are assumed to be measurable without explicit reference. Alternately, we also view X and Y as random variables on some other, unspecified, probability space; e.g., when they are viewed as real-valued measurements. In this case P is viewed as a joint distribution on X and Y . In either case, the probability of a set $T \subset Z$ is defined by

$$P(T) = \int_T dP(z)$$

(where $z = (x, y)$ with $x \in X$ and $y \in Y$) and the expectation of function f on Z is denoted by

$$E(f) = \int_Z f(z) dP(z).$$

When Z is countable we also use P , with some abuse of notation, for the probability mass function; i.e., for $z \in Z$, $P(z)$ denotes $P(\{z\})$. Hence $P(T) = \sum_{z \in T} P(z)$ and $E(f) = \sum_{z \in Z} f(z) P(z)$ in this case. When Z is continuous, a density associated with P (if it exists) is denoted by p .

When Z is countable we use $P(y|x)$ to denote the probability that $Y = y$ given that $X = x$ (viewing X and Y as random variables) and similarly for $P(x|y)$. Hence $P(\cdot|x)$ denotes the conditional distribution on Y , given $X = x$. The marginal distribution in X is defined by⁷ $P_{|x}(x) = \sum_{y \in Y} P(x, y)$. Here and elsewhere, we abbreviate $P((x, y))$ by $P(x, y)$.

⁶ If Z is countable then we assume this σ -algebra contains all subsets of Z ; otherwise we assume that Z is a complete, separable metric space (see Section 9.1) and that this σ -algebra is the smallest σ -algebra that contains the open sets of Z (i.e., the σ -algebra of Borel sets).

⁷ When Z is uncountable, the marginal and conditional distributions are defined so that

$$\int_Z f(x, y) dP(x, y) = \int_X \left(\int_Y f(x, y) dP(y|x) \right) dP_{|x}(x)$$

for every bounded measurable function f .

Finally, we list some other notation that is used several places in the text, indicating which section it is defined in:

X, Y, A, \mathcal{H} , and l	Sections 1.1 and 2.1
\mathcal{P}	Section 2.1
$\mathbf{r}_{h,l}(P), \mathbf{r}_h(P)$ (true risk)	Section 2.2
$\mathbf{r}_l^*(P), \mathbf{r}^*(P)$ (optimal risk)	Section 2.2
$\hat{\mathbf{r}}_h(\tilde{z})$ (empirical risk)	Section 2.2
$\hat{\mathbf{r}}^*(\tilde{z})$ (optimal empirical risk)	Section 2.2
d_v	Section 2.2
$L, L_\varepsilon, L_{\alpha,v}$ (regret functions)	Section 2.3
R (big ‘L’ risk)	Section 2.3
$m(\varepsilon, \delta), m(\alpha, v, \delta)$ (sample complexity)	Section 2.4
\mathcal{N} (covering number)	Sections 10.1 and 3.2
\mathcal{M} (packing number)	Section 10.1
dim (metric dimension)	Section 10.1
dim_P (pseudo dimension)	Section 4
\mathcal{C} (capacity)	Section 6
ρ_l	Section 6
$l_{\mathcal{H}}$	Section 3
$\mathbf{F}_{ \tilde{z}}$	Section 3
$\hat{\mathbf{E}}$ (empirical expectation)	Section 3
d_{L^1} (L^1 distance for vectors)	Section 3.2
$d_{L^1(P)}$ (L^1 distance for functions)	Section 4
$d_{L^1(P,\rho)}$ (L^1 distance for functions)	Section 6

2. LEARNING AND OPTIMIZATION

We now further formalize the basic problem of learning, as introduced in Section 1.1. We introduce a formal notion of a learning algorithm, and a higher level loss function, which we call a *regret function*, that measures how well the learning algorithm performs. The regret function is defined in terms of the low level loss function l discussed in the previous section. Finally, we show how an algorithm can solve the learning problem by solving a related optimization problem.

2.1. The Basic Components $X, Y, A, \mathcal{H}, \mathcal{P}$, and l

We first review and further formalize the six components of the basic learning problem introduced in the previous section: $X, Y, A, \mathcal{H}, \mathcal{P}$, and l .

The first four components are the instance, outcome, decision, and decision rule spaces, respectively. The first three of these are arbitrary sets, and the fourth, \mathcal{H} , is a family of functions from X into A . These have been discussed extensively in the previous section.

The fifth component, \mathcal{P} , is a family of joint probability distributions on $X \times Y$. These represent the possible “states of nature” that might govern the generation of examples. The set $Z = X \times Y$ is called the *sample space*. We assume that examples are drawn independently at random according to some probability distribution $P \in \mathcal{P}$ on the sample space Z . A sequence of examples is called a *sample*. In what follows⁸ we usually assume that \mathcal{P} includes all probability distributions on Z . Hence our results are distribution independent.

The last component, the loss function l , is a mapping from $Y \times A$ into \mathbb{R} . In this paper we assume that l is bounded and nonnegative; i.e., $0 \leq l \leq M$ for some real M . When Y and A are finite it is always possible to enforce this condition by simply adding a constant to l , which does not change the learning problem in any essential way. When either Y or A is infinite, the learning problem sometimes needs to be restricted to meet this condition. For example, in regression⁹ we might restrict the possible parameter vectors in A and/or the possible outcomes in Y such that for every $y \in Y$ and $a \in A$, $\hat{P}(y; a) \geq b$ for some constant b . We can then take $M = -\log b$. In density estimation, the same thing can be accomplished by restricting the instance space X to a bounded subset of \mathbb{R}^n on which all densities in \mathcal{H} have values uniformly greater than b and less than B for constants $0 < b < B$. We can then add $\log B$ to the loss function to make it positive. The same method works for estimating distributions on discrete spaces: we restrict ourselves to a finite instance space X and demand that for all $x \in X$ and all probability distributions $h \in \mathcal{H}$, $h(x) \geq b > 0$ (see, e.g., Abe and Warmuth, 1990; Yamanishi, 1990). These restrictions are often reasonable in practice—e.g., most measurements naturally have bounded ranges—but

⁸ It is, however, possible and in fact common to assume that \mathcal{P} is a very specific class of probability distributions on Z . For example, let $X = \mathbb{R}^n$. Then if we are doing classification learning and Y is discrete we may assume that y is selected according to an arbitrary distribution on Y , and for each y , $P(x|y)$ is a multi-variate Gaussian distribution on X (Duda and Hart, 1973). On the other hand, if we are doing linear regression, then Y is real-valued and we might assume that x is selected according to an arbitrary distribution on X , and y is a linear function of x with additive Gaussian noise. In PAC learning theory we have a discrete analog of the latter case. Here we usually have $X = \{0, 1\}^n$, $Y = \{0, 1\}$, and y a Boolean function of x of a particular type (e.g., defined by a small disjunctive normal form formula), possibly plus random noise.

⁹ Note that to get bounded loss in linear regression, X must be a bounded subset of \mathbb{R}^n as well, since we can't bound Y without bounding X . The coefficients of the functions in \mathcal{H} must also be bounded.

they can be annoying (see Vapnik (1989) and Pollard (1984, 1990) for alternative approaches for unbounded loss functions).

2.2. Measuring Distance from Optimality with the d_v Metric

For a given decision rule $h \in \mathcal{H}$ and distribution P on the sample space Z , the expected loss of h is the average value of $l(y, h(x))$, where the example (x, y) is drawn at random according to P . It is defined by

$$\mathbf{r}_{h,l}(P) = \mathbf{r}_h(P) = \mathbf{E}(l(y, h(x))) = \int_Z l(y, h(x)) dP(x, y)$$

(the subscript l is omitted when the loss function is clear from the context). Since l is bounded, this expectation is finite for every distribution P . In decision theory the expected loss $\mathbf{r}_h(P)$ is called the *risk* of h when P is the true underlying distribution. This quantity generalizes the notion of the *error* of h used in computational learning theory.

In Section 1.1 we stated the goal of learning quite informally: Given examples chosen independently at random from some unknown probability distribution $P \in \mathcal{P}$, find a decision rule \hat{h} in \mathcal{H} that comes “close to” minimizing the risk $\mathbf{r}_h(P)$ over all $h \in \mathcal{H}$. Let $\mathbf{r}_l^*(P)$ (or $\mathbf{r}^*(P)$ when l is clear from the context) denote the infimum of $\mathbf{r}_h(P)$ over all h in the decision rule space \mathcal{H} . To formalize our notion of a basic learning problem, we first need to say what we mean that $\mathbf{r}_{\hat{h}}(P)$ is “close to” $\mathbf{r}^*(P)$.

Let $r = \mathbf{r}_{\hat{h}}(P)$ and $s = \mathbf{r}^*(P)$. One natural interpretation is to demand that $|r - s| \leq \varepsilon$ for some small $\varepsilon > 0$. However, we see in Section 3.1 that sometimes it is better to use a relative measure of distance. For any real $v > 0$, let d_v be the function defined by

$$d_v(r, s) = \frac{|r - s|}{v + r + s}$$

for any non-negative reals r and s . It is straightforward but tedious to verify that d_v is a metric on \mathfrak{R}^+ . The d_v metric is similar to the standard function

$$\frac{|r - s|}{s}$$

used to measure the difference between the quality r of a given solution and the quality s of an optimal solution in combinatorial optimization. However, our measure has been modified to be well-behaved when one or

both of its arguments are zero, and to be symmetric in its arguments (so that it is a metric). Three other properties of d_v are also useful:

1. For all non-negative reals r and s , $0 \leq d_v(r, s) < 1$.
2. For all non-negative $r \leq s \leq t$, $d_v(r, s) \leq d_v(r, t)$ and $d_v(s, t) \leq d_v(r, t)$.
3. For $0 \leq r, s \leq M$, $|r - s|/(v + 2M) \leq d_v(r, s) \leq |r - s|/v$.

We refer to the second property by saying that d_v is *compatible with the ordering on the reals*.

2.3. The Regret Function L and the Big “ L ” Risk R

Once we have specified how we measure closeness to optimality, we still need to specify our criteria for a successful learning algorithm. Do we need to have the risk of the decision rule found close to the optimum $\mathbf{r}_I^*(P)$ with high probability, or should its average distance from $\mathbf{r}_I^*(P)$ be small? Do we measure success in terms of the performance of the algorithm on the worst case distribution in \mathcal{P} , or do we use some average case analysis over distributions in \mathcal{P} ? These questions lead us right back to decision theory again, but this time at a higher level in the analysis of learning.

To see this, consider the structure of a learning algorithm \mathcal{A} . For any sample size m , the algorithm \mathcal{A} may be given a sample $\vec{z} = ((x_1, y_1), \dots, (x_m, y_m))$ drawn at random from Z^m according to an unknown product distribution P^m , where $P \in \mathcal{P}$. For any such \vec{z} it chooses a decision rule $\mathcal{A}(\vec{z}) \in \mathcal{H}$. Thus abstractly, the algorithm defines a function \mathcal{A} from the set of all samples over Z into \mathcal{H} ; i.e., $\mathcal{A}: \bigcup_{m \geq 1} Z^m \rightarrow \mathcal{H}$. Since we are not requiring computability here, we call such \mathcal{A} a *learning method*. When $P \in \mathcal{P}$ is the actual “state of nature” governing the generation of examples, and the algorithm produces the decision rule $h \in \mathcal{H}$, let us say that we suffer a non-negative real-valued *regret* $L(P, h)$. Thus, formally $L: \mathcal{P} \times \mathcal{H} \rightarrow \mathbb{R}^+$. In our treatment here, the regret function L is derived from the loss function l , and measures the extent to which we have failed to produce a near optimal decision rule, assuming P is the true state of nature (i.e., the amount of “regret” we feel for not having produced the optimal decision rule). Finally, for each possible state of nature P , the average regret suffered by the algorithm, over all possible training samples $\vec{z} \in Z^m$, is the *big “ L ” risk* of that algorithm under P for sample size m . This big “ L ” risk is defined formally by

$$R_{L, \mathcal{A}, m}(P) = \int_{\vec{z} \in Z^m} L(P, \mathcal{A}(\vec{z})) dP^m(\vec{z}).$$

The goal of learning is to minimize big “ L ” risk.

We illustrate these definitions with a few examples. First suppose we want to capture the notion of successful learning that is used in the PAC model. Then one possibility is to introduce an *accuracy parameter* $\varepsilon > 0$ and define the regret function $L = L_\varepsilon$ by letting $L_\varepsilon(P, h) = 1$ if $\mathbf{r}_{h,l}(P) - \mathbf{r}_l^*(P) > \varepsilon$, and $L_\varepsilon(P, h) = 0$ otherwise. Thus we suffer regret only when the decision rule h produced by the learning algorithm has risk that is more than ε from optimal, measured by the absolute difference metric. For this definition of regret, the big “L” risk $R_{L_\varepsilon, \mathcal{A}, m}(P)$ measures the probability that the decision rule produced by \mathcal{A} has risk more than ε from optimal, when \mathcal{A} is given m random training examples drawn according to P . We then demand that this big “L” risk be small, i.e., smaller than some given *confidence parameter* $\delta > 0$.

In the PAC model it is commonly assumed that the examples given to the algorithm \mathcal{A} are noise-free examples of some underlying target function $f \in \mathcal{H}$. In this case the risk $\mathbf{r}^*(P)$ of the optimal decision rule in \mathcal{H} is zero, and hence $L_\varepsilon(P, h) = 1 \Leftrightarrow \mathbf{r}_{h,l}(P) > \varepsilon$. Hence demanding big “L” risk at most δ gives the usual PAC criterion that the risk (or “error”) of the decision rule (or “hypothesis”) produced by \mathcal{A} must be greater than ε with probability at most δ .

The regret function L can also be defined similarly, but using the d_v metric to measure distance from optimality, instead of the absolute difference. Specifically, for every $v > 0$ and $0 < \alpha < 1$, we can define the regret function $L_{\alpha, v}$ by letting $L_{\alpha, v}(P, h) = 1$ if $d_v(\mathbf{r}_{h,l}(P), \mathbf{r}_l^*(P)) > \alpha$, and $L_{\alpha, v}(P, h) = 0$ otherwise. In this case the big “L” risk $R_{L_{\alpha, v}, \mathcal{A}, m}(P)$ measures the probability that the risk of the decision rule produced by the algorithm \mathcal{A} has distance more than α from optimal in the d_v metric, when the algorithm is given m random training examples drawn according to P . We see in Sections 2.4 and 3.1 why this sometimes gives a more useful and flexible definition of regret. In fact, in this paper, we will give our main results in terms of the family $\{L_{\alpha, v} : v > 0 \text{ and } 0 < \alpha < 1\}$ of regret functions, and show how corresponding results may be derived as corollaries for the family $\{L_\varepsilon : \varepsilon > 0\}$ of regret functions.

Other regret functions are also possible and lead to different learning criteria. For example, another, perhaps simpler, way to define regret is to let $L(P, h) = \mathbf{r}_{h,l}(P) - \mathbf{r}_l^*(P)$. When $\mathbf{r}_l^*(P) = 0$, as it does in the standard noise-free PAC model, this definition makes the regret L equal to the risk $\mathbf{r}_{h,l}(P)$; i.e., the expectation of the underlying loss l . In this case the big “L” risk $R_{L, \mathcal{A}, m}(P)$ measures the expectation of the loss incurred by the learning algorithm \mathcal{A} when it is given m random training examples drawn according to P , forms a decision rule h , and then uses h to determine the action on one further independent random example drawn according to P . This gives a generalization of the learning criterion studied in Haussler *et al.* (1990). When $\mathbf{r}_l^*(P) \neq 0$, the big “L” risk gives the expectation of the

amount of such loss above and beyond the expected loss that would be suffered if the optimal decision rule were used. In particular, in density estimation, where P and h are both densities on the instance space X , if $P \in \mathcal{H}$ then defining the regret by $L(P, h) = r_{h,l}(P) - r_l^*(P)$ makes it equal to the Kullback–Leibler divergence from P to h . Hence the big “L” risk is the expected Kullback–Leibler divergence of the decision rule h returned by the algorithm from the true density (see Sections 1.1.3 and 1.1.4).

It is also possible to define the regret function L directly, without using an underlying loss function l . For example, in density estimation it is possible to use other measures of the distance between two densities; e.g., the Hellinger distance or the total variational distance, as in Barron and Cover (1990) and Yamanishi (1990). The criterion from Kearns and Schapire (1990) for inferring a good *model of probability* can also be defined using an appropriate regret function, without defining an underlying loss l .

2.4. Full Formalization of the Basic Learning Problem

Having defined the regret function, and thereby the big “L” risk function, we still face one last issue: do we want to minimize big “L” risk in the worst case over all possible states of nature P in \mathcal{P} , or do we want to assume a *prior distribution* on possible distributions in \mathcal{P} , so that we can define a notion of “average case” big “L” risk to be minimized? The former goal is known as *minimax optimality*, and has been used in the PAC model. The latter is the *Bayesian* notion of optimality (Berger, 1985; Kiefer, 1987), and has been used in several approaches to learning in neural nets based on statistical mechanics (Denker *et al.* 1987; Tishby *et al.*, 1989; Gyorgyi and Tishby, 1990; Sompolinsky *et al.*, 1990; Oppen and Haussler, 1991a, 1991b). Unfortunately this last question has no clear cut answer, and leads us directly into a longstanding unresolved debate in statistics (see, e.g., Lindley (1990) and following discussion.). Since we have set out to generalize the PAC model, and since our results are best illustrated in the minimax setting, we formalize the notion of a basic learning problem using the minimax criterion. In subsequent work we hope to further explore this Bayesian setting. (For recent work in Bayesian approaches to neural network learning see MacKay (1992), Buntine and Weigend (1991), and for Bayesian versions of the PAC model see Haussler *et al.* (1991b) and Buntine (1990).)

We can now define exactly what we mean by a basic learning problem, and what it means for a learning method to solve this problem in this minimax setting.

DEFINITION 1. A basic learning problem is defined by six components, X , Y , A , \mathcal{H} , \mathcal{P} , and \mathcal{L} , where the first five components are as defined in

Section 2.1, and the last component, \mathcal{L} , is a family of regret functions as defined in Section 2.3 (e.g., $\mathcal{L} = \{L_{\alpha, \nu} : \nu > 0 \text{ and } 0 < \alpha < 1\}$, or $\mathcal{L} = \{L_\varepsilon : \varepsilon > 0\}$ for some loss function l). Let \mathcal{A} be a learning method as defined in Section 2.3. We say that \mathcal{A} solves the basic learning problem if for all $L \in \mathcal{L}$ and all $0 < \delta < 1$ there exists a finite sample size $m = m(L, \delta)$ such that

$$\text{for all } P \in \mathcal{P}, \quad R_{L, \mathcal{A}, m}(P) \leq \delta.$$

The *sample complexity* of the learning method \mathcal{A} is the smallest such integer valued function $m(L, \delta)$. When $\mathcal{L} = \{L_{\alpha, \nu} : \nu > 0 \text{ and } 0 < \alpha < 1\}$ we denote $m(L_{\alpha, \nu}, \delta)$ by $m(\alpha, \nu, \delta)$ and when $\mathcal{L} = \{L_\varepsilon : \varepsilon > 0\}$ we denote $m(L_\varepsilon, \delta)$ by $m(\varepsilon, \delta)$.

As discussed above, this definition generalizes the PAC criterion, and several others as well. In fact, this definition is quite generous, in that sample size needed to get the big “L” risk less than δ is only required to be finite for each $\delta > 0$. In particular, using property (3) of the d_ν metric from Section 2.2, when the underlying loss function l is bounded, as we assume here, any algorithm \mathcal{A} solves the basic learning problem using the $L_{\alpha, \nu}$ class of regret functions if and only if it solves it using the L_ε class. Thus it does not matter which of these two classes of regret functions we use. However, in practice it is the sample complexity of \mathcal{A} that is critical, and this will depend on which class of regret functions are used.

The nature of this dependence is seen more clearly when we expand the condition

$$R_{L, \mathcal{A}, m}(P) \leq \delta$$

for $L = L_\varepsilon$ and $L = L_{\alpha, \nu}$. When $L = L_\varepsilon$, this condition means that given m random training examples drawn according to P , with probability at least $1 - \delta$, the decision rule \hat{h} produced by the algorithm \mathcal{A} satisfies

$$\mathbf{r}_{\hat{h}, l}(P) \leq \mathbf{r}_l^*(P) + \varepsilon;$$

i.e., the risk of \hat{h} is at most ε greater than that of the optimal decision rule in \mathcal{H} . When $L = L_{\alpha, \nu}$, this condition is the same, except that we require

$$\mathbf{r}_{\hat{h}, l}(P) \leq \frac{1 + \alpha}{1 - \alpha} \mathbf{r}_l^*(P) + \frac{\alpha \nu}{1 - \alpha}.$$

Thus in the former case, the sample complexity is defined in terms of small additive deviations from optimality, and in the latter, we allow both additive and multiplicative deviations. These deviations are controlled by the parameters α and ν .

For example, when $\mathbf{r}_l^*(P) = 0$ as in the standard PAC model, then setting $\alpha = \frac{1}{2}$ and $v = \varepsilon$ makes the L_ε and $L_{\alpha,v}$ conditions equivalent; each reduces to the PAC condition

$$\mathbf{r}_{h,l}(P) \leq \varepsilon.$$

When $\mathbf{r}_l^*(P) > 0$, the $L_{\alpha,v}$ condition approximates the L_ε condition when α is small and $v \approx \varepsilon/\alpha$. In particular, since we are assuming that the underlying loss function l is bounded between 0 and M , we have $0 \leq \mathbf{r}_{h,l}(P)$, $\mathbf{r}_l^*(P) \leq M$, and property (3) of the d_v metric shows that the $L_{\alpha,v}$ condition with $v = 2M$ and $\alpha = \varepsilon/4M$ implies the L_ε condition. This shows how the two parameter $L_{\alpha,v}$ condition is generally more flexible than the single parameter L_ε condition.

2.5. Relation between Learning and Optimization

Let us assume that the underlying loss function l is fixed, and we are using either the L_ε or the $L_{\alpha,v}$ regret functions derived from l . In order to solve a basic learning problem, we must find, with high probability, a decision rule \hat{h} with risk close to optimal. As the true distribution P is unknown, to do this we must rely on estimates of $\mathbf{r}_h(P)$ for the various $h \in \mathcal{H}$ which are derived from the given random training sample. For a given $h \in \mathcal{H}$ and training sample $\vec{z} = (z_1, \dots, z_m)$, where $z_i = (x_i, y_i) \in Z$, let $\hat{\mathbf{r}}_h(\vec{z})$ denote the empirical risk on \vec{z} ; i.e., $\hat{\mathbf{r}}_h(\vec{z}) = (1/m) \sum_{i=1}^m l(y_i, h(x_i))$. Let $\hat{\mathbf{r}}^*(\vec{z}) = \inf\{\hat{\mathbf{r}}_h(\vec{z}) : h \in \mathcal{H}\}$. We can then define a natural optimization problem associated with the basic learning problem: given the training sample \vec{z} , find a decision rule $\hat{h} \in \mathcal{H}$ such that $\hat{\mathbf{r}}_{\hat{h}}(\vec{z})$ is close to $\hat{\mathbf{r}}^*(\vec{z})$, i.e., a decision rule whose empirical risk on the training sample is close to minimal.

Solving the optimization problem does not automatically solve the learning problem. We need to have good empirical risk estimates as well. Since l is bounded, for every $h \in \mathcal{H}$, as the sample size $m \rightarrow \infty$, $\hat{\mathbf{r}}_h(\vec{z}) \rightarrow \mathbf{r}_h(P)$ with probability 1. We will say that the empirical risk estimates of decision rules in \mathcal{H} converge *uniformly* to the true risk if for all ε and $\delta > 0$ there exists a sample size m such that when the $z_i \in \vec{z}$, $1 \leq i \leq m$, are drawn independently at random from Z according to the distribution P , with probability at least $1 - \delta$, we have $\rho(\hat{\mathbf{r}}_h(\vec{z}), \mathbf{r}_h(P)) \leq \varepsilon$ for all $h \in \mathcal{H}$. Here ρ is some metric on \mathfrak{R}^+ ; e.g., either the absolute difference or the d_v metric.

The following result shows that uniform convergence of the empirical risk estimates, along with a learning method \mathcal{A} that gives a randomized solution to the optimization problem on the estimates, gives a solution to the basic learning problem. We state it for the d_v metric, but the same argument works also for the absolute difference metric.

LEMMA 1. Let $v > 0$ and $0 < \alpha, \delta < 1$. Suppose the sample size $m = m(\alpha, v, \delta)$ is such that for all probability distributions $P \in \mathcal{P}$

$$\Pr(\exists h \in \mathcal{H} : d_v(\hat{r}_h(\vec{z}), r_h(P)) > \alpha/3) \leq \delta/2,$$

where the $z_i \in \bar{Z}$, $1 \leq i \leq m$, are drawn independently at random from Z according to the distribution P . Suppose also that the algorithm \mathcal{A} is such that for all $P \in \mathcal{P}$

$$\Pr(d_v(\hat{r}_{\mathcal{A}(\vec{z})}(\vec{z}), \hat{r}^*(\vec{z})) > \alpha/3) \leq \delta/2,$$

where \vec{z} is drawn randomly by P as above. Then for all $P \in \mathcal{P}$

$$\Pr(d_v(r_{\mathcal{A}(\vec{z})}(P), r^*(P)) > \alpha) \leq \delta;$$

i.e., \mathcal{A} solves the basic learning problem for the family of $L_{\alpha, v}$ regret functions and has sample complexity at most $m(\alpha, v, \delta)$.

Proof. By the triangle inequality for d_v , if

1. $d_v(r_{\mathcal{A}(\vec{z})}(P), \hat{r}_{\mathcal{A}(\vec{z})}(\vec{z})) \leq \alpha/3$,
2. $d_v(\hat{r}_{\mathcal{A}(\vec{z})}(\vec{z}), \hat{r}^*(\vec{z})) \leq \alpha/3$, and
3. $d_v(\hat{r}^*(\vec{z}), r^*(P)) \leq \alpha/3$,

then

$$d_v(r_{\mathcal{A}(\vec{z})}(P), r^*(P)) \leq \alpha.$$

The second assumption of the lemma states that (2) holds with probability at least $1 - \delta/2$. The first assumption implies that both (1) and (3) hold with probability at least $1 - \delta/2$. (If (3) fails then we can find a decision rule $h \in \mathcal{H}$ such that $d_v(\hat{r}_h(\vec{z}), r_h(P)) > \alpha/3$. Here we use the compatibility of d_v with the ordering on the reals.) Hence with probability at least $1 - \delta$ all of (1)–(3) hold. The result follows. ■

In statistics, this type of result is called a *consistency theorem* about the “statistic” (i.e., the decision rule) computed by the learning method \mathcal{A} . This use of the term “consistency” differs sharply from that common in PAC learning research.

3. UNIFORMLY GOOD EMPIRICAL ESTIMATES OF MEANS

In this section we concentrate on the problem of bounding the number of random examples needed to get good empirical estimates of the risk of each of the decision rules in a decision rule space \mathcal{H} . For each decision rule

$h \in \mathcal{H}$ and example $z = (x, y) \in Z$, let $l_h(z) = l(y, h(x))$. As in the previous section, we assume that l is a nonnegative bounded loss function taking values in the interval $[0, M]$; thus for each decision rule h , l_h defines a random variable taking values in $[0, M]$. The value of l_h on an example (x, y) is the loss incurred when h is used to determine the action to take, for instance x , and the outcome is y . The risk of h is just the expectation of l_h ; i.e.,

$$r_h(P) = \mathbf{E}(l_h) = \int_Z l_h(z) dP(z).$$

Furthermore, if $\vec{z} = (z_1, \dots, z_m)$ is a sequence of examples from Z , then the empirical risk of h on \vec{z} is the empirical estimate of the mean of l_h based on the sample \vec{z} , which we denote by $\hat{\mathbf{E}}_{\vec{z}}(l_h)$; i.e.,

$$\hat{r}_h(\vec{z}) = \hat{\mathbf{E}}_{\vec{z}}(l_h) = \frac{1}{m} \sum_{i=1}^m l_h(z_i).$$

Let $l_{\mathcal{H}} = \{l_h : h \in \mathcal{H}\}$. We need to draw enough random examples to get a uniformly good empirical estimate of the expectation of every random variable in $l_{\mathcal{H}}$.

The general problem of obtaining a uniformly good estimate of the expectation of every function in a class \mathbf{F} of real-valued functions has been widely studied (see, e.g., Vapnik, 1982, Pollard, 1984, Dudley, 1984, and their references). If no assumptions at all are made about the functions in \mathbf{F} , we immediately run into the problem that some functions in \mathbf{F} could take on arbitrarily large values with arbitrarily small probabilities, making it impossible to obtain uniformly good empirical estimates of all expectations with any finite sample size. This problem can be avoided by making assumptions about the moments of the functions in \mathbf{F} , as in Vapnik (1982), or by assuming that there exists a single non-negative function with a finite expectation (called an *envelope*) that lies above the absolute value of every function in \mathbf{F} , as in Pollard (1984) and Dudley (1984). In our case, when the loss takes only values in $[0, M]$, then the constant function M serves as an envelope. This case is especially nice since this same envelope works for all distributions on the domain Z of the functions in \mathbf{F} .

The usual measure of deviation of empirical estimates from true means is simply the absolute value of the difference. Thus we would say that the empirical estimates for the expectations of the functions in \mathbf{F} converge uniformly to the true expectations if as the size of the random sample \vec{z} grows,

$$\Pr(\exists f \in \mathbf{F}: |\hat{\mathbf{E}}_{\vec{z}}(f) - \mathbf{E}(f)| > \varepsilon)$$

goes to zero for any $\varepsilon > 0$. (This is called (*uniform*) *convergence in probability*; see, e.g., Billingsley (1986)). Vapnik, Dudley, Pollard, and others have obtained general bounds on the sample size needed so that

$$\Pr(\exists f \in \mathbf{F}: |\hat{\mathbf{E}}_z(f) - \mathbf{E}(f)| > \varepsilon) < \delta$$

for $\varepsilon, \delta > 0$ (Pollard, 1984; Dudley, 1984; Vapnik, 1982). Vapnik also obtains better bounds in some important cases by considering the relative deviation of empirical estimates from true expectations. He looks at bounds on the sample size needed so that

$$\Pr\left(\exists f \in \mathbf{F}: \frac{\mathbf{E}(f) - \hat{\mathbf{E}}_z(f)}{\mathbf{E}(f)} > \varepsilon\right) < \delta$$

and also bounds on the sample size needed so that

$$\Pr\left(\exists f \in \mathbf{F}: \frac{\mathbf{E}(f) - \hat{\mathbf{E}}_z(f)}{\sqrt{\mathbf{E}(f)}} > \varepsilon\right) < \delta.$$

(Anthony and Shawe-Taylor (1990) also obtain bounds of the latter form.) Note that these are one-sided bounds, in that they only bound the probability that the empirical mean is significantly smaller than the true mean. While extremely useful, as we mentioned in the previous section, these measures of deviation suffer from a discontinuity at $\mathbf{E}(f) = 0$, and lack of convenient metric properties. As in Pollard (1986), we give bounds on the sample size needed so that

$$\begin{aligned} & \Pr(\exists f \in \mathbf{F}: d_v(\hat{\mathbf{E}}_z(f), \mathbf{E}(f)) > \alpha) \\ &= \Pr\left(\exists f \in \mathbf{F}: \frac{|\hat{\mathbf{E}}_z(f) - \mathbf{E}(f)|}{v + \hat{\mathbf{E}}_z(f) + \mathbf{E}(f)} > \alpha\right) < \delta, \end{aligned}$$

i.e., the deviation measured using the d_v metric.¹⁰ By setting v and α appropriately, we obtain results similar to those of Pollard (1984) and Vapnik (1982) as special cases of our main theorem. However, our results are restricted to the case where all functions in \mathbf{F} are positive and uniformly bounded.

¹⁰ Pollard (1986) also gives results that can be used to bound the sample size needed so that

$$\Pr\left(\exists f \in \mathbf{F}: \frac{|\hat{\mathbf{E}}_z(f) - \mathbf{E}(f)|}{v + \sqrt{\hat{\mathbf{E}}_z(f)} + \sqrt{\mathbf{E}(f)}} > \alpha\right) < \delta,$$

in analogy with the second type of bound given by Vapnik, except that these bounds are two-sided. We do not pursue this further here.

3.1. The Case of Finite F

Before we consider the general case, it is useful to see what bounds we can get in the case that F is a finite set of functions. Here we can easily prove the following.

THEOREM 1. *Let F be a finite set of functions on Z with $0 \leq f(z) \leq M$ for all $f \in F$ and $z \in Z$. Let $\vec{z} = (z_1, \dots, z_m)$ be a sequence of m examples drawn independently from Z according to any distribution on Z , and let $\varepsilon > 0$. Then*

$$\Pr(\exists f \in F: |\hat{E}_{\vec{z}}(f) - E(f)| > \varepsilon) \leq 2|F| e^{-2\varepsilon^2 m/M^2}.$$

For $0 < \delta \leq 1$ and sample size

$$m \geq \frac{M^2}{2\varepsilon^2} \left(\ln |F| + \ln \frac{2}{\delta} \right)$$

this probability is at most δ . Further, for any $v > 0$ and $0 < \alpha < 1$,

$$\Pr(\exists f \in F: d_v(\hat{E}_{\vec{z}}(f), E(f)) > \alpha) \leq 2|F| e^{-\alpha^2 v m/M}.$$

For $0 < \delta \leq 1$ and sample size

$$m \geq \frac{M}{\alpha^2 v} \left(\ln |F| + \ln \frac{2}{\delta} \right)$$

this probability is at most δ .

Proof. For the second part of the theorem, using Bernstein's inequality (see, e.g., Pollard, 1984) it is easy to show that for any single function f with $0 \leq f \leq M$,

$$\Pr(d_v(\hat{E}_{\vec{z}}(f), E(f)) > \alpha) < 2e^{-\alpha^2 v m/M}.$$

Details are given in Lemma 9, part (2) in the Appendix. It follows that the probability that there is any $f \in F$ with $d_v(\hat{E}_{\vec{z}}(f), E(f)) > \alpha$ is at most $2|F| e^{-\alpha^2 v m/M}$. Setting this bound to δ and solving for m gives the result on the sample size. The proof of the first part of the lemma is similar, except we use Hoeffding's inequality (see, e.g., Pollard, 1984), which implies that for any single f ,

$$\Pr(|\hat{E}_{\vec{z}}(f) - E(f)| > \varepsilon) \leq 2e^{-2\varepsilon^2 m/M^2}. \quad \blacksquare$$

By letting $F = I_{\mathcal{X}}$, this theorem can be used in conjunction with Lemma 1 from the previous section to obtain bounds on the sample complexity of learning algorithms that minimize empirical risk. Here we can

use either the L_ε or the $L_{\alpha,v}$ family of regret functions. In the former case we get a sample complexity

$$m(\varepsilon, \delta) = O\left(\frac{M^2}{\varepsilon^2} \left(\log |I_{\mathcal{H}}| + \log \frac{1}{\delta}\right)\right). \quad (1)$$

In the latter case we get a sample complexity

$$m(\alpha, v, \delta) = O\left(\frac{M}{\alpha^2 v} \left(\log |I_{\mathcal{H}}| + \log \frac{1}{\delta}\right)\right). \quad (2)$$

As shown in the previous section, a generalization of the PAC learning model can be obtained by using either the L_ε or the $L_{\alpha,v}$ regret functions, in the latter case by setting $\alpha = \frac{1}{2}$ and $v = \varepsilon$. Note that plugging this latter setting into (2) gives a sample complexity

$$m(\varepsilon, \delta) = O\left(\frac{M}{\varepsilon} \left(\log |I_{\mathcal{H}}| + \log \frac{1}{\delta}\right)\right), \quad (3)$$

a significant improvement over (1), which is quadratic in M/ε . Thus the generalization of the PAC model using the d_v metric to measure distance from optimality, and the resulting $L_{\alpha,v}$ family of regret functions, offers new insight in this regard. (Vapnik's (1982) use of the relative difference between empirical estimates and true expectations also has this advantage; see Anthony and Shawe-Taylor (1990) and also the Appendix of Blumer *et al.* (1989).)

3.2. The General Case

The main task of this section is to generalize Theorem 1 to infinite collections of uniformly bounded functions. The basic idea is simple: we replace the infinite class F of functions with a finite class F_0 that "approximates" it, in the sense that each function in F is close to some function in F_0 , and argue that some type of uniform convergence of empirical estimates for F_0 implies uniform convergence for F . In the simplest version of this technique, the choice of F_0 depends only on F and the distribution P , as in the "direct method" discussed in Section II.2 of Pollard (1984) (see also Vapnik (1982, Sect. 6.6), Dudley (1984, Chap. 6), Benedek and Itai (1988), and White (1990a)). However, more general results (apart from certain measurability constraints) are obtained by allowing F_0 to depend on the particular random sample \vec{z} (e.g., Pollard (1984, Chap. 2)). Here F_0 is called a "random cover," and its size is called a "random covering number." It is this type of result that we derive here.

We need a few preliminary definitions to introduce the notions of ε -covers and metric dimension. A more general treatment of these ideas is given in the Appendix, Section 9.1. This more general treatment will be used later, but the following definitions suffice for this section.

For any real vectors $\vec{x} = (x_1, \dots, x_m)$ and $\vec{y} = (y_1, \dots, y_m)$ in \mathfrak{R}^m , let $d_{L^1}(\vec{x}, \vec{y}) = (1/m) \sum_{i=1}^m |x_i - y_i|$. Thus d_{L^1} is the L^1 distance metric. Let T be a set of points that lie in a bounded region of \mathfrak{R}^m . For any $\varepsilon > 0$, an ε -cover for T is a finite set $N \subset \mathfrak{R}^m$ (not necessarily contained in T) such that for all $\vec{x} \in T$ there is a $\vec{y} \in N$ with $d_{L^1}(\vec{x}, \vec{y}) \leq \varepsilon$. The function $\mathcal{N}(\varepsilon, T)$ denotes the size of the smallest ε -cover for T . We refer to $\mathcal{N}(\varepsilon, T)$ as a *covering number*.

Following Kolmogorov and Tihomirov (1961) we define the *upper metric dimension* of the set T of points by

$$\overline{\dim}(T) = \limsup_{\varepsilon \rightarrow 0} \frac{\log \mathcal{N}(\varepsilon, T)}{\log(1/\varepsilon)}.$$

The *lower metric dimension*, denoted by $\underline{\dim}$, is defined similarly using \liminf . When $\overline{\dim}(T) = \underline{\dim}(T)$, then this quantity is denoted $\dim(T)$, and referred to simply as the *metric dimension* of T . Note that if $\mathcal{N}(\varepsilon, T) = (g(\varepsilon)/\varepsilon)^n$, where $g(\varepsilon)$ is polylogarithmic in $1/\varepsilon$, then $\dim(T) = n$. Hence the metric dimension essentially picks out the exponent in the rate of growth of the covering number as a function of $1/\varepsilon$.

Assume all functions in F map from Z into $[0, M]$. For any sample $\vec{z} = (z_1, \dots, z_m)$, with $z_i \in Z$, let

$$F_{|\vec{z}} = \{(f(z_1), \dots, f(z_m)): f \in F\}.$$

We call $F_{|\vec{z}}$ the *restriction* of F to \vec{z} . Note that $F_{|\vec{z}}$ is a set of points in the m -cube $[0, M]^m$. We can consider the size of the covering number $\mathcal{N}(\varepsilon, F_{|\vec{z}})$ as giving some indication of the “richness at scale $\approx \varepsilon$ ” of the class F of functions, restricted to the domain z_1, \dots, z_m . The metric dimension of $F_{|\vec{z}}$ gives some indication of the “number of essential degrees of freedom” in this restriction of F .

When z_1, \dots, z_m are drawn independently at random from Z , the *random covering number* $E(\mathcal{N}(\varepsilon, F_{|\vec{z}}))$ gives some indication of the “richness” of F on a “typical” set of m points in the domain Z . Note that for finite F , we have $\mathcal{N}(\varepsilon, F_{|\vec{z}}) \leq |F|$ for all ε and all samples \vec{z} , and hence the random covering number $E(\mathcal{N}(\varepsilon, F_{|\vec{z}})) \leq |F|$ for all ε , all sample sizes m , and all distributions on Z . The main result about uniform empirical estimates for infinite classes of functions is similar to Theorem 1 except that the random covering numbers are used in place of $|F|$.

THEOREM 2 (Pollard, 1986). *Let \mathbf{F} be a permissible¹¹ set of functions on Z with $0 \leq f(z) \leq M$ for all $f \in \mathbf{F}$ and $z \in Z$. Let $\vec{z} = (z_1, \dots, z_m)$ be a sequence of m examples drawn independently from Z according to any distribution on Z . Then for any $v > 0$ and $0 < \alpha < 1$,*

$$\Pr(\exists f \in \mathbf{F}: d_v(\hat{\mathbf{E}}_{\vec{z}}(f), \mathbf{E}(f)) > \alpha) \leq 4\mathbf{E}(\mathcal{N}(\alpha v/8, \mathbf{F}_{|\vec{z}})) e^{-\alpha^2 v m/16M}.$$

COROLLARY 1 (Pollard, 1984). *Under the same assumptions as above, for all $\varepsilon > 0$,*

$$\Pr(\exists f \in \mathbf{F}: |\hat{\mathbf{E}}_{\vec{z}}(f) - \mathbf{E}(f)| > \varepsilon) \leq 4\mathbf{E}(\mathcal{N}(\varepsilon/16, \mathbf{F}_{|\vec{z}})) e^{-\varepsilon^2 m/128M^2}.$$

Proof of Corollary. This follows directly from the above result by setting $v = 2M$, and $\alpha = \varepsilon/4M$. To see this, note that property (3) of the d_v metric (section 2.2) implies that $|r - s| \leq \varepsilon$ whenever $d_v(r, s) \leq \alpha$ for all $0 \leq r, s \leq M$ when this setting of v and α is used. ■

The constants in these results are only crude estimates. No serious attempt has been made to minimize them. (See the recent results of Talagrand (1991) for much better constants for Corollary 1).

The bound in this latter result depends critically on the relative magnitudes of the negative exponent in $e^{-\varepsilon^2 m/128M^2}$ and the exponent in the expectation of the covering number $\mathcal{N}(\varepsilon/16, \mathbf{F}_{|\vec{z}})$, which reflects the extent to which $\mathbf{F}_{|\vec{z}}$ “fills up” the m -cube $[0, M]^m$. For example, if $\mathbf{F}_{|\vec{z}}$ has metric dimension at most n for all m and all \vec{z} , then there is a constant c_0 such that for any $\eta > 0$, $\mathcal{N}(\varepsilon/16, \mathbf{F}_{|\vec{z}}) \leq (c_0 M/\varepsilon)^{n+\eta}$ for suitably small ε . In this case the negative exponential term eventually dominates the expected covering number, and beyond a critical sample size

$$m_0 = O\left(\frac{nM^2}{\varepsilon^2} \log \frac{M}{\varepsilon}\right),$$

the bound goes to zero exponentially fast. We see examples of this in the following section, where we give bounds on the metric dimension of $\mathbf{F}_{|\vec{z}}$ in terms of a combinatorial parameter called the pseudo dimension of \mathbf{F} . The theorem actually shows that this exponential drop off occurs even if this metric dimension bound holds only for “most” \vec{z} .

On the other hand, if with high probability $\mathbf{F}_{|\vec{z}}$ “fills up” the m -cube $[0, M]^m$ to the extent that $\mathcal{N}(\varepsilon/16, \mathbf{F}_{|\vec{z}}) \approx (c_0/\varepsilon)^m$, which is as large as possible, then the covering number dominates, and the bound is trivial. Results in Vapnik (1982), (Theorem A.2, p. 220) indicate that uniform convergence does not take place in this case. Similar remarks apply to the bound given in Theorem 2, which uses the d_v metric.

¹¹ This is a measurability condition defined in Pollard (1984) which need not concern us in practice. Further details are given in Section 9.2 of the Appendix.

The proof of Theorem 2 follows the proof of Pollard's Theorem 24 (Pollard, 1984, p. 25) in general outline. However, the use of the d_v metric necessitates a number of substantial modifications. The approach taken here is different from that taken (independently, but prior to this work) in Pollard (1986). Still different, and more involved, techniques are used in the more general theory of weighted empirical processes developed by Alexander (1985, 1987). While the proof of Theorem 2 we give is simpler, it is still somewhat lengthy, so it is given in Appendix 9.4.

Actually, we can prove a slightly stronger result than Theorem 2. This result is obtained by bounding the probability of uniform convergence on a sample of length m in terms of the expected covering numbers associated with a sample of length $2m$, and by expanding the expectation to include the negative exponential term with a "truncation" at 1. It turns out that this saves us a factor of $\frac{1}{2}$ in the negative exponential term. We also include special bounds for the case that $\mathbf{F}_{|\bar{z}}$ is always finite. This case comes up, for example, when $\mathbf{F} = l_{\mathcal{X}}$ and we use the discrete loss function l , as in the PAC learning model.

THEOREM 3. *Let \mathbf{F} be a permissible set of functions on Z with $0 \leq f(z) \leq M$ for all $f \in \mathbf{F}$ and $z \in Z$. Assume $v > 0$, $0 < \alpha < 1$, and $m \geq 1$. Suppose that \bar{z} is generated by m independent random draws according to any probability measure on Z . Let*

$$p(\alpha, v, m) = \mathbf{Pr}\{\bar{z} \in Z^m: \exists f \in \mathbf{F} \text{ with } d_v(\hat{\mathbf{E}}_{\bar{z}}(f), \mathbf{E}(f)) > \alpha\}.$$

Then

$$p(\alpha, v, m) \leq 2\mathbf{E}(\min(2\mathcal{N}(\alpha v/8, \mathbf{F}_{|\bar{z}}, d_{L^1}) e^{-\alpha^2 v m/8M}, 1)),$$

where the expectation is over \bar{z} drawn randomly from Z^{2m} . If in addition $\mathbf{F}_{|\bar{z}}$ is finite for all $\bar{z} \in Z^{2m}$ then

$$p(\alpha, v, m) \leq 2\mathbf{E}(\min(2|\mathbf{F}_{|\bar{z}}| e^{-\alpha^2 v m/2M}, 1)).$$

Theorem 2 is obtained as a corollary of this result by substituting $m/2$ for m and not taking the minimum with 1 in the left hand side of the first bound for $p(\alpha, v, m)$. We use Theorem 3 to obtain slightly better constants in some of the results in the sequel.

4. PSEUDO DIMENSION OF CLASSES OF REAL-VALUED FUNCTIONS

In this section we look at one way that bounds on the covering numbers appearing in Theorem 2 can be obtained. This technique, due to Pollard (1984), who extended methods from Dudley (1978), is based on certain

intuitions from combinatorial geometry. It generalizes the techniques based on the Vapnik–Chervonenkis dimension used in Blumer *et al.* (1989), which apply only to $\{0, 1\}$ -valued functions. We begin by establishing some basic notation.

DEFINITION 2. For $x \in \mathfrak{R}$, let $\text{sign}(x) = 1$ if $x > 0$ else $\text{sign}(x) = 0$. For $\vec{x} = (x_1, \dots, x_d) \in \mathfrak{R}^d$, let $\text{sign}(\vec{x}) = (\text{sign}(x_1), \dots, \text{sign}(x_d))$ and for $T \subset \mathfrak{R}^d$ let $\text{sign}(T) = \{\text{sign}(\vec{x}) : \vec{x} \in T\}$. For any Boolean vector $\vec{b} = (b_1, \dots, b_d)$, $\{\vec{x} \in \mathfrak{R}^d : \text{sign}(\vec{x}) = \vec{b}\}$ is called the \vec{b} -orthant of \mathfrak{R}^d , where we have, somewhat arbitrarily, included points with value zero for a particular coordinate in the associated lower orthant. Thus $\text{sign}(T)$ denotes the set of orthants intersected by T . For any $T \subset \mathfrak{R}^d$, and $\vec{x} \in \mathfrak{R}^d$, let $T + \vec{x} = \{\vec{y} + \vec{x} : \vec{y} \in T\}$; i.e., the translation of T obtained by adding the vector \vec{x} . We say that T is *full* if there exists $\vec{x} \in \mathfrak{R}^d$ such that $\text{sign}(T + \vec{x}) = \{0, 1\}^d$; i.e., if there exists some translation of T that intersects all 2^d orthants of \mathfrak{R}^d .

The following result is well known and can be proved in a variety of ways. For example, it follows easily from well known bounds on the number of cells in arrangements of hyperplanes (see, e.g., Edelsbrunner, 1987). We give an elementary proof using a technique from Dudley (1978).

LEMMA 2. *No hyperplane in \mathfrak{R}^d intersects all orthants of \mathfrak{R}^d .*

Proof. Let T be a hyperplane in \mathfrak{R}^d . Choose a vector $\vec{x} \in \mathfrak{R}^d$ as follows. If T includes the origin, then let \vec{x} be any vector that is orthogonal to T and has at least one strictly negative coordinate. (For any non-zero orthogonal vector \vec{x} , if \vec{x} does not have a negative coordinate then $-\vec{x}$ does.) Otherwise let \vec{x} be the (non-zero) vector in T on the line perpendicular to T that passes through the origin. To complete the proof, we show that for all $\vec{y} \in T$, $\text{sign}(\vec{y}) \neq \vec{1} - \text{sign}(\vec{x})$, where $\vec{1}$ denotes the all 1's vector.

Suppose to the contrary that $\text{sign}(\vec{y}) = \vec{1} - \text{sign}(\vec{x})$ for some $\vec{y} \in T$. This implies that the inner product $\sum_{i=1}^d x_i y_i$ is non-positive, and is in fact strictly negative if either \vec{x} or \vec{y} contain a strictly negative coordinate. However, by our choice of \vec{x} , either \vec{x} is orthogonal to \vec{y} and contains a strictly negative coordinate, giving an immediate contradiction, or \vec{x} is non-zero and \vec{x} is orthogonal to $\vec{y} - \vec{x}$. In this last case,

$$\sum_{i=1}^d x_i y_i = \sum_{i=1}^d x_i^2,$$

which is again a contradiction, since the left side is non-positive while the right side is strictly positive. ■

It follows from this lemma that if T is contained in a hyperplane of \mathfrak{R}^d then T is not full.

DEFINITION 3. Let \mathbf{F} be a family of functions from a set Z into \mathfrak{R} . For any sequence $\vec{z} = (z_1, \dots, z_d)$ of points in Z , let $\mathbf{F}_{|\vec{z}} = \{(f(z_1), \dots, f(z_d)) : f \in \mathbf{F}\}$. If $\mathbf{F}_{|\vec{z}}$ is full then we say that \vec{z} is *shattered* by \mathbf{F} . The *pseudo dimension* of \mathbf{F} , denoted $\mathbf{dim}_p(\mathbf{F})$, is the largest d such that there exists a sequence of d points in Z that is shattered by \mathbf{F} . If arbitrarily long finite sequences are shattered, then $\mathbf{dim}_p(\mathbf{F})$ is finite.

It is clear that when \mathbf{F} is a set of $\{0, 1\}$ -valued functions then for any sequence \vec{z} of d points in Z , $\mathbf{F}_{|\vec{z}}$ is full if and only if $\mathbf{F}_{|\vec{z}} = \{0, 1\}^d$. Thus in this case $\mathbf{dim}_p(\mathbf{F})$ is the length d of the longest sequence of points \vec{z} such that $\mathbf{F}_{|\vec{z}} = \{0, 1\}^d$. This is the definition of the Vapnik–Chervonenkis dimension of a class \mathbf{F} of $\{0, 1\}$ -valued functions (Vapnik, 1982; Haussler and Welzl, 1987; Blumer *et al.*, 1989). Thus the pseudo dimension generalizes the Vapnik–Chervonenkis dimension to arbitrary classes of real-valued functions.

The pseudo dimension also generalizes the algebraic notion of the dimension of a vector space of real-valued functions (Dudley, 1978).

THEOREM 4 (Dudley). *Let \mathbf{F} be a d -dimensional vector space of functions from a set Z into \mathfrak{R} . Then $\mathbf{dim}_p(\mathbf{F}) = d$.*

Proof. Fix any sequence $\vec{z} = (z_1, \dots, z_{d+1})$ of points in Z . For any $f \in \mathbf{F}$ let $\Psi(f) = (f(z_1), \dots, f(z_{d+1}))$. Then Ψ is a linear mapping from \mathbf{F} into \mathfrak{R}^{d+1} , and the image of Ψ is $\mathbf{F}_{|\vec{z}}$. Since \mathbf{F} is a vector space of dimension d , this implies that $\mathbf{F}_{|\vec{z}}$ is a subspace of \mathfrak{R}^{d+1} of dimension at most d . Hence by Lemma 2, $\mathbf{F}_{|\vec{z}}$ is not full. This implies $\mathbf{dim}_p(\mathbf{F}) \leq d$. On the other hand, if \mathbf{F} is a d -dimensional vector space of real-valued functions on Z , then there exists a sequence \vec{z} of d points in Z such that $\mathbf{F}_{|\vec{z}} = \mathfrak{R}^d$. Hence \vec{z} is shattered, implying that $\mathbf{dim}_p(\mathbf{F}) \geq d$. ■

There are many other ways that the VC dimension can be generalized to real-valued functions (Natarajan, 1989a, 1989b; Pollard, 1984; Vapnik, 1989; Dudley, 1987). Dudley (1987) compares several such generalizations, albeit in a different context. The generalization we have proposed here, the pseudo dimension, is a minor variant of the notion used by Pollard (1984) to define classes of real-valued functions of polynomial discrimination, called VC-subgraph classes in Dudley (1987). The pseudo dimension will be used in the form defined above in Pollard's new book (Pollard, 1990).

The pseudo dimension has a few invariance properties that are useful (see Pollard, 1990, for further results of this type).

THEOREM 5. *Let \mathbf{F} be a family of functions from Z into \mathfrak{R} . Fix any function g from Z into \mathfrak{R} and let $\mathbf{G} = \{g + f : f \in \mathbf{F}\}$. Let I be a real interval (possibly all of \mathfrak{R}) such that every function in \mathbf{F} takes values only in I .*

Fix any nondecreasing (resp. nonincreasing) function $h: I \rightarrow \mathfrak{R}$ and let $\mathbf{H} = \{h \circ f: f \in \mathbf{F}\}$, where \circ indicates function composition. Then

1 (Wenocur and Dudley, 1981). $\dim_{\mathbf{P}}(\mathbf{G}) = \dim_{\mathbf{P}}(\mathbf{F})$ and

2 (Nolan and Pollard, 1987; Dudley, 1987). $\dim_{\mathbf{P}}(\mathbf{H}) \leq \dim_{\mathbf{P}}(\mathbf{F})$, with equality if h is continuous and strictly increasing (resp. continuous and strictly decreasing).

Proof. Part (1) follows directly from the fact that the notion of a set of points being full is invariant under translation. For part (2) it suffices to prove the results for h nondecreasing and h continuous and strictly increasing. Let $\vec{z} = (z_1, \dots, z_d)$ be such that $\mathbf{H}_{|\vec{z}}$ is full, i.e., such that $\mathbf{H}_{|\vec{z}} - \vec{x}$ intersects all 2^d orthants of \mathfrak{R}^d for some vector $\vec{x} = (x_1, \dots, x_d)$ in \mathfrak{R}^d . Then for every Boolean vector $\vec{b} \in \{0, 1\}^d$ there exists a function $f_{\vec{b}} \in \mathbf{F}$ such that for every i , $1 \leq i \leq d$, we have $h \circ f_{\vec{b}}(z_i) > x_i$ if and only if the i th bit of \vec{b} is 1. For each i , $1 \leq i \leq d$, let

$$u_i = \min\{f_{\vec{b}}(z_i): \text{the } i\text{th bit of } \vec{b} \text{ is } 1\}$$

and

$$l_i = \max\{f_{\vec{b}}(z_i): \text{the } i\text{th bit of } \vec{b} \text{ is } 0\}.$$

Since h is nondecreasing, we have $u_i > l_i$ for each i . Let $r_i = (u_i + l_i)/2$ for each i and $\vec{r} = (r_1, \dots, r_d)$. Let $T = \{f_{\vec{b}}: \vec{b} \in \{0, 1\}^d\}$. Then clearly $T_{|\vec{z}} - \vec{r}$ intersects every orthant of \mathfrak{R}^d , so $T_{|\vec{z}}$ is full. Since $T \subset \mathbf{F}$, this implies that $\mathbf{F}_{|\vec{z}}$ is full, and hence $\dim_{\mathbf{P}}(\mathbf{H}) \leq \dim_{\mathbf{P}}(\mathbf{F})$. Equality follows when h is continuous and strictly increasing since we obtain the class \mathbf{F} from \mathbf{H} by composing with h^{-1} . ■

By putting a probability measure on Z , we can view a class \mathbf{F} of real-valued functions on Z as a pseudo metric space. The distance between two functions is the integral of the absolute value of their difference, i.e., the L^1 distance, relative to the given measure. To make this work, we need to make some assumptions about the integrability of the functions in \mathbf{F} under the given measure. Since we will be concerned only with families of functions taking values in a bounded range in this paper, this will cause no problems for us. For convenience, we choose this range to be $[0, M]$. For a more general treatment, see Pollard (1984) and Dudley (1984).

DEFINITION 4. Let \mathbf{F} be a class of functions from Z into $[0, M]$, where $M > 0$, and let P be a probability measure on Z . Then $d_{L^1(P)}$ is the pseudo metric on \mathbf{F} defined by

$$d_{L^1(P)}(f, g) = \mathbf{E}(|f - g|) = \int_Z |f(z) - g(z)| dP(z) \quad \text{for all } f, g \in \mathbf{F}.$$

The notions of ε -cover and metric dimension used in the previous section can be generalized to arbitrary pseudo metric spaces. This generalization is given in Section 9.1 of the Appendix. In the remainder of the paper we use the concepts and notation given there without further special reference.

Using techniques that go back to Dudley (1978), Pollard has obtained a beautiful theorem bounding the metric dimension of $(\mathbf{F}, d_{L^1(P)})$ by $\dim_P(\mathbf{F})$ for any probability measure P on Z . Actually this result is much stronger in that it gives explicit bounds on the packing number for \mathbf{F} using $d_{L^1(P)}$ balls of radius ε . Since packing numbers are closely related to covering numbers (Theorem 12 in Section 9.1), these bounds can then be used with Theorem 2 to obtain uniform convergence results for empirical estimates of functions in \mathbf{F} . We now state and prove a version of Pollard's result (Pollard, 1984, Lemma 25, p. 27) for the special case when \mathbf{F} is a class of functions taking values in the interval $[0, M]$ with somewhat better bounds on the packing numbers.

THEOREM 6 (Pollard). *Let \mathbf{F} be a family of functions from a set Z into $[0, M]$, where $\dim_P(\mathbf{F}) = d$ for some $1 \leq d < \infty$. Let P be a probability measure on Z . Then for all $0 < \varepsilon \leq M$,*

$$\mathcal{M}(\varepsilon, \mathbf{F}, d_{L^1(P)}) < 2 \left(\frac{2eM}{\varepsilon} \ln \frac{2eM}{\varepsilon} \right)^d.$$

The proof we give uses essentially the same techniques as Pollard's, with some minor modifications. It relies on a few lemmas, which we give now. The first, which we give without proof, was discovered independently by a number of people (see Assouad, 1983) including Vapnik and Chervonenkis (1971), but is most often attributed to Sauer (1972) in the computer science literature.

LEMMA 3 (Sauer). *Let \mathbf{F} be a class of functions from $S = \{1, 2, \dots, m\}$ into $\{0, 1\}$ with $|\mathbf{F}| > 1$ and let d be the length of the longest sequence of points \vec{z} from S such that $F_{|\vec{z}} = \{0, 1\}^d$. Then¹²*

$$|\mathbf{F}| \leq \sum_{i=0}^d \binom{m}{i} \leq (em/d)^d,$$

where e is the base of the natural logarithm. ■

¹² See, e.g., Dudley (1984, Proposition 2.2.9) or Blumer *et al.* (1989, Appendix), for a proof of the second inequality. Note also that Vapnik and Chervonenkis (1971) actually contains a slightly weaker result.

In the next lemma we bound the packing numbers of $(\mathbf{F}, d_{L^1(P)})$ in terms of the expected number of orthants intersected by a random translation of a random restriction of \mathbf{F} . This is the key lemma of the proof.

LEMMA 4. *Let \mathbf{F} be a family of functions from a set Z into $[0, M]$ and let P be a probability measure on Z . Let $\vec{r} = (r_1, \dots, r_m)$ be a random vector in $[0, M]^m$, where each r_i is drawn independently at random from the uniform distribution on $[0, M]$. Let $\vec{z} = (z_1, \dots, z_m)$ be a random vector in Z^m , where each z_i is drawn independently at random from P . Then for all $\varepsilon > 0$*

$$\mathbf{E}(|\text{sign}(\mathbf{F}_{|\vec{z}} - \vec{r})|) \geq \mathcal{M}(\varepsilon, \mathbf{F}, d_{L^1(P)})(1 - \mathcal{M}(\varepsilon, \mathbf{F}, d_{L^1(P)})e^{-\varepsilon m/M}).$$

Proof. For all $f \in \mathbf{F}$ we denote $(f(z_1), \dots, f(z_m))$ by $f_{|\vec{z}}$. Choose $\varepsilon > 0$. Let \mathbf{G} be an ε -separated subset of \mathbf{F} (w.r.t. $d_{L^1(P)}$), with $|\mathbf{G}| = \mathcal{M}(\varepsilon, \mathbf{F}, d_{L^1(P)})$. Then

$$\begin{aligned} & \mathbf{E}(|\text{sign}(\mathbf{F}_{|\vec{z}} - \vec{r})|) \\ & \geq \mathbf{E}(|\text{sign}(\mathbf{G}_{|\vec{z}} - \vec{r})|) \\ & \geq \mathbf{E}(|\{f \in \mathbf{G}: \text{sign}(f_{|\vec{z}} - \vec{r}) \neq \text{sign}(g_{|\vec{z}} - \vec{r}) \quad \text{for all } g \in \mathbf{G}, g \neq f\}|)|) \\ & = \sum_{f \in \mathbf{G}} \Pr(\text{sign}(f_{|\vec{z}} - \vec{r}) \neq \text{sign}(g_{|\vec{z}} - \vec{r}) \quad \text{for all } g \in \mathbf{G}, g \neq f) \\ & = \sum_{f \in \mathbf{G}} (1 - \Pr(\exists g \in \mathbf{G}, g \neq f: \text{sign}(f_{|\vec{z}} - \vec{r}) = \text{sign}(g_{|\vec{z}} - \vec{r}))) \\ & \geq \sum_{f \in \mathbf{G}} (1 - |\mathbf{G}| \max_{g \in \mathbf{G}, g \neq f} \Pr(\text{sign}(f_{|\vec{z}} - \vec{r}) = \text{sign}(g_{|\vec{z}} - \vec{r}))). \end{aligned}$$

Let f and g be distinct functions in \mathbf{G} . Since \mathbf{G} is ε -separated,

$$\int_Z |f(z) - g(z)| dP(z) > \varepsilon.$$

In addition, the range of f and g is $[0, M]$. Hence if z_i is drawn at random from P and r_i is drawn at random from the uniform distribution on $[0, M]$, then the probability that r_i lies between $f(z_i)$ and $g(z_i)$ is at least ε/M . And $\text{sign}(f_{|\vec{z}} - \vec{r}) = \text{sign}(g_{|\vec{z}} - \vec{r})$ only if this fails to occur for each i , $1 \leq i \leq m$. Thus

$$\Pr(\text{sign}(f_{|\vec{z}} - \vec{r}) = \text{sign}(g_{|\vec{z}} - \vec{r})) \leq \left(1 - \frac{\varepsilon}{M}\right)^m \leq e^{-\varepsilon m/M}.$$

Since this holds for every distinct f and g in \mathbf{G} , combining this with the inequality above, we have

$$\mathbf{E}(|\text{sign}(\mathbf{F}_{|\bar{z}} - \bar{r})|) \geq |\mathbf{G}|(1 - |\mathbf{G}|e^{-\epsilon m/M}).$$

Since $|\mathbf{G}| = \mathcal{M}(\epsilon, \mathbf{F}, d_{L^1(P)})$, this gives the result. ■

Proof of Theorem 6. Since $\dim_P(\mathbf{F}) = d$, it follows from Sauer's lemma that $|\text{sign}(\mathbf{F}_{|\bar{z}} - \bar{r})| \leq (\epsilon m/d)^d$ for all $m \geq d$, $\bar{z} \in Z^m$ and $\bar{r} \in [0, M]^m$. Hence the above lemma implies that

$$(\epsilon m/d)^d \geq \mathcal{M}(\epsilon, \mathbf{F}, d_{L^1(P)})(1 - \mathcal{M}(\epsilon, \mathbf{F}, d_{L^1(P)})e^{-\epsilon m/M}) \quad (4)$$

for all probability measures P on Z and $m \geq d$. It is easily verified that if

$$\frac{M}{\epsilon} \ln(2\mathcal{M}(\epsilon, \mathbf{F}, d_{L^1(P)})) < d$$

then the upper bound given in Theorem 6 follows trivially using the fact that $\epsilon \leq M$. Thus we may assume that $(M/\epsilon) \ln(2\mathcal{M}(\epsilon, \mathbf{F}, d_{L^1(P)})) \geq d$. Hence, if $m \geq (M/\epsilon) \ln(2\mathcal{M}(\epsilon, \mathbf{F}, d_{L^1(P)}))$, then $m \geq d$ and

$$(1 - \mathcal{M}(\epsilon, \mathbf{F}, d_{L^1(P)})e^{-\epsilon m/M}) \geq \frac{1}{2}.$$

Thus from (4) we obtain

$$\left(\frac{eM \ln(2\mathcal{M}(\epsilon, \mathbf{F}, d_{L^1(P)}))}{\epsilon d} \right)^d \geq \frac{1}{2} \mathcal{M}(\epsilon, \mathbf{F}, d_{L^1(P)}).$$

With some simple calculations, this gives the bound of Theorem 6. ■

Using our results on uniform convergence from Sections 3.2 and 9.4, we can now show the following.

THEOREM 7. *Let \mathbf{F} be a permissible family of functions from a set Z into $[0, M]$ with $\dim_P(\mathbf{F}) = d$ for some $1 \leq d < \infty$. Assume $m \geq 1$, $0 < v \leq 8M$, and $0 < \alpha < 1$. Let \bar{z} be generated by m independent draws according to any distribution on Z . Then*

$$\Pr(\exists f \in \mathbf{F}: d_v(\hat{\mathbf{E}}_{\bar{z}}(f), \mathbf{E}(f)) > \alpha) \leq 8 \left(\frac{16eM}{\alpha v} \ln \frac{16eM}{\alpha v} \right)^d e^{-\alpha^2 v m / 8M}.$$

Moreover, for $m \geq (8M/\alpha^2 v)(2d \ln(8eM/\alpha v) + \ln(8/\delta))$ this probability is at most δ .

Proof. Let $\varepsilon = \alpha v/8$. Since $\alpha < 1$ and $v \leq 8M$, $\varepsilon \leq M$. For any sequence \vec{z} of points in Z there is a trivial isometry between $(\mathbf{F}_{|\vec{z}}, d_{L^1})$ and $(\mathbf{F}, d_{L^1(P_{\vec{z}})})$, where $P_{\vec{z}}$ is the empirical measure induced by \vec{z} , in which each set has measure equal to the fraction of the points in \vec{z} it contains. Thus by Theorem 12 of Section 9.1 and Theorem 6, we have

$$\mathcal{N}(\varepsilon, \mathbf{F}_{|\vec{z}}, d_{L^1}) \leq \mathcal{M}(\varepsilon, \mathbf{F}_{|\vec{z}}, d_{L^1}) \leq 2 \left(\frac{2eM}{\varepsilon} \ln \frac{2eM}{\varepsilon} \right)^d,$$

for all $\vec{z} \in Z^{2m}$. Hence the given probability is at most

$$8 \left(\frac{2eM}{\varepsilon} \ln \frac{2eM}{\varepsilon} \right)^d e^{-\alpha^2 v m / 8M} = 8 \left(\frac{16eM}{\alpha v} \ln \frac{16eM}{\alpha v} \right)^d e^{-\alpha^2 v m / 8M}$$

by Theorem 3.

For the second result, setting the bound above equal to δ and solving for m gives

$$m \geq \frac{8M}{\alpha^2 v} \left(d \ln \left(\frac{16eM}{\alpha v} \ln \frac{16eM}{\alpha v} \right) + \ln \frac{8}{\delta} \right).$$

It is easily verified that $\ln(a \ln a) < 2 \ln(a/2)$ when $a \geq 5$, and from this the bound given in the second result follows. ■

COROLLARY 2. *Under the same assumptions as above, for all $0 < \varepsilon \leq M$,*

$$\Pr(\exists f \in \mathbf{F}: |\hat{\mathbf{E}}_{|\vec{z}}(f) - \mathbf{E}(f)| > \varepsilon) \leq 8 \left(\frac{32eM}{\varepsilon} \ln \frac{32eM}{\varepsilon} \right)^d e^{-\varepsilon^2 m / 64M^2}.$$

Moreover, for $m \geq (64M^2/\varepsilon^2)(2d \ln(16eM/\varepsilon) + \ln(8/\delta))$ this probability is at most δ .

Proof. This follows directly from the above result by setting $v = 2M$, $\alpha = \varepsilon/4M$, and using property (3) of the d_v metric, as in the proof of Corollary 1 in Section 3.2. ■

5. SOME APPLICATIONS OF PSEUDO DIMENSION IN LEARNING

We now look at how the theoretical results obtained in the previous two sections can be applied to certain types of learning problems. Suppose that we have a basic learning problem defined by X , Y , A , \mathcal{H} , \mathcal{P} , and \mathcal{L} , where \mathcal{L} is the family of $L_{x,v}$ regret functions for an underlying loss function l , as in Section 2. As before, let $Z = X \times Y$, $l_h: Z \rightarrow [0, M]$ be defined by $l_h(x, y) = l(y, h(x))$ for all $h \in \mathcal{H}$, and $l_{\mathcal{H}} = \{l_h: h \in \mathcal{H}\}$. In this section we

show how to obtain sample complexity bounds on algorithms for this basic learning problem using Theorem 7 above.

To obtain these bounds, we need bounds on $\dim_{\mathbf{P}}(l_{\mathcal{H}})$. In this section we look at some useful tricks for computing $\dim_{\mathbf{P}}(l_{\mathcal{H}})$ in the important case $A \subset \mathfrak{R}$, i.e., when each decision is represented by a real number. In the following section we discuss more general decision spaces.

When $A \subset \mathfrak{R}$, the functions in \mathcal{H} are themselves real-valued, so we can talk about the pseudo dimension of \mathcal{H} itself, without reference to any particular loss function. What makes this useful is that in many important cases the pseudo dimension of \mathcal{H} is the same as the pseudo dimension of $l_{\mathcal{H}}$. Thus we can factor out the effects of the loss function in deriving our sample size bounds, and concentrate on the pseudo dimension of the decision rule space \mathcal{H} .

DEFINITION 5. Let $l: Y \times A \rightarrow \mathfrak{R}$ be a loss function, where $A \subset \mathfrak{R}$ and Y is an arbitrary set. For each $y \in Y$ define the function $f_y: A \rightarrow \mathfrak{R}$ by letting $f_y(a) = l(y, a)$ for each $a \in A$; i.e., f_y is the restriction of l obtained by fixing its argument to y . We say that l is *monotone* over A if for every $y \in Y$, either f_y is strictly increasing on A , or f_y is strictly decreasing on A . Thus f_y may be increasing for some $y \in Y$ and decreasing for others.

LEMMA 5. If $A \subset \mathfrak{R}$ and l is a loss function on $A \times Y$ that is monotone over A , then $\dim_{\mathbf{P}}(l_{\mathcal{H}}) = \dim_{\mathbf{P}}(\mathcal{H})$.

Proof. Suppose that $\vec{x} = (x_1, \dots, x_d)$ is shattered by \mathcal{H} , where $x_i \in X$, $1 \leq i \leq d$. Then there is some real vector $\vec{r} = (r_1, \dots, r_d)$ such that $\mathcal{H}_{|\vec{x}} - \vec{r}$ intersects all 2^d orthants of \mathfrak{R}^d . Hence for every Boolean vector $\vec{b} \in \{0, 1\}^d$ there exists a function $h_{\vec{b}} \in \mathcal{H}$ such that for every i , $1 \leq i \leq d$, we have $h_{\vec{b}}(x_i) > r_i$ if and only if the i th bit of \vec{b} is 1. Fix an outcome $y \in Y$. Let $\vec{z} = (z_1, \dots, z_d)$, where $z_i = (x_i, y)$ for all i , $1 \leq i \leq d$. Note that if f_y is strictly increasing, then for any $h \in \mathcal{H}$ and $1 \leq i \leq d$, $h(x_i) > r_i \Leftrightarrow l_h(z_i) = l(y, h(x_i)) = f_y(h(x_i)) > f_y(r_i)$. Hence, for every Boolean vector $\vec{b} \in \{0, 1\}^d$ there exists a function $h_{\vec{b}} \in \mathcal{H}$ such that for every i , $1 \leq i \leq d$, we have $l_{h_{\vec{b}}}(z_i) > f_y(r_i)$ if and only if the i th bit of \vec{b} is 1. A similar result holds if f_y is strictly decreasing. Thus \vec{z} is shattered by $l_{\mathcal{H}}$. It follows that $\dim_{\mathbf{P}}(l_{\mathcal{H}}) \geq \dim_{\mathbf{P}}(\mathcal{H})$.

For the other direction, assume $\vec{z} = (z_1, \dots, z_d)$ is shattered by $l_{\mathcal{H}}$, where $z_i = (x_i, y_i)$ for all i , $1 \leq i \leq d$. We show that $\vec{x} = (x_1, \dots, x_d)$ is shattered by \mathcal{H} . Since \vec{z} is shattered by $l_{\mathcal{H}}$, there is some real vector $\vec{r} = (r_1, \dots, r_d)$ such that for every Boolean vector $\vec{b} \in \{0, 1\}^d$ there exists a function $h_{\vec{b}} \in \mathcal{H}$ such that for every i , $1 \leq i \leq d$, we have $l_{h_{\vec{b}}}(z_i) > r_i$ if and only if the i th bit of \vec{b} is 1. Let $A_0 = \{h_{\vec{b}}(x_i): 1 \leq i \leq d \text{ and } \vec{b} \in \{0, 1\}^d\}$. For each outcome $y \in Y$

define the function $g_y: \mathfrak{R} \rightarrow A_0$ as follows. If f_y is increasing then $g_y(r) = \max\{a \in A_0: f_y(a) \leq r\}$ and if f_y is decreasing then $g_y(r) = \max\{a \in A_0: f_y(a) > r\}$. Then for each i , $1 \leq i \leq d$, we either have

1. for all $h \in \mathcal{H}$, $l_h(z_i) = f_{y_i}(h(x_i)) > r_i \Leftrightarrow h(x_i) > g_{y_i}(r_i)$ or
2. for all $h \in \mathcal{H}$, $l_h(z_i) = f_{y_i}(h(x_i)) > r_i \Leftrightarrow h(x_i) \leq g_{y_i}(r_i)$.

Hence, for every Boolean vector $\vec{b} \in \{0, 1\}^d$ there exists a function $h'_b \in \mathcal{H}$ such that for every i , $1 \leq i \leq d$, we have $h'_b(x_i) > g_{y_i}(r_i)$ if and only if the i th bit of \vec{b} is 1. (To see this, let \vec{c} be the Boolean vector derived from \vec{b} by complementing the bit in each position i for which f_{y_i} is decreasing, and then let $h'_b = h_{\vec{c}}$.) Thus $\vec{x} = (x_1, \dots, x_d)$ is shattered by \mathcal{H} . It follows that $\dim_P(\mathcal{H}) \geq \dim_P(l_{\mathcal{H}})$, and combined with the above inequality, this gives the result. ■

Combined with Theorem 7, this gives the following result on the uniform convergence of empirical risk estimates for the basic learning problem. Here and below we use the notation $\hat{\mathbf{r}}_{h,i}(\vec{z})$ and $\mathbf{r}_{h,i}(P)$ introduced in Section 2 for the empirical risk estimate and true risk of a decision rule h , respectively.

THEOREM 8. *Assume the decision space $A \subset \mathfrak{R}$, the loss function l is monotone over A and bounded between 0 and M , the decision rule space \mathcal{H} is such that $l_{\mathcal{H}}$ is permissible, and $1 \leq d = \dim_P(\mathcal{H}) < \infty$. Assume $m \geq 1$, $0 < v < 8M$, and $0 < \alpha < 1$. Let P be any probability distribution on Z . Let \vec{z} be generated by m independent draws from Z according to P . Then*

$$\Pr(\exists h \in \mathcal{H}: d_v(\hat{\mathbf{r}}_{h,i}(\vec{z}), \mathbf{r}_{h,i}(P)) > \alpha) \leq 8 \left(\frac{16eM}{\alpha v} \ln \frac{16eM}{\alpha v} \right)^d e^{-\alpha^2 v m / 8M}.$$

Moreover, for $m \geq (8M/\alpha^2 v)(2d \ln(8eM/\alpha v) + \ln(8/\delta))$ this probability is at most δ .

Proof. By Lemma 5, $\dim_P(l_{\mathcal{H}}) = \dim_P(\mathcal{H})$ in this case. Thus, since $\hat{\mathbf{r}}_{h,i}(\vec{z}) = \hat{\mathbf{E}}_{\vec{z}}(l_h)$ and $\mathbf{r}_{h,i}(P) = \mathbf{E}(l_h)$, the result follows directly from Theorem 7. ■

When its conditions are satisfied, this theorem, combined with Lemma 1 from Section 2.5, gives us a means of bounding in terms of $\dim_P(\mathcal{H})$ the sample complexity $m(\alpha, v, \delta)$ of any algorithm that solves the basic learning problem by returning (with high probability) a decision rule with near minimal empirical risk on the training sample. The resulting bound is

$$m(\alpha, v, \delta) = O \left(\frac{M}{\alpha^2 v} \left(\dim_P(\mathcal{H}) \log \frac{M}{\alpha v} + \log \frac{1}{\delta} \right) \right). \quad (5)$$

This is similar to the sample complexity

$$m(\alpha, v, \delta) = O\left(\frac{M}{\alpha^2 v} \left(\log |\mathcal{H}| + \log \frac{1}{\delta}\right)\right)$$

that can be obtained by using Theorem 1, when \mathcal{H} (and hence $l_{\mathcal{H}}$) is finite. The term $\mathbf{dim}_P(\mathcal{H}) \log(M/\alpha v)$ replaces the term $\log |\mathcal{H}|$. In particular, for the “PAC settings” $\alpha = \frac{1}{2}$ and $v = \varepsilon$ we get the sample complexity

$$m(\varepsilon, \delta) = O\left(\frac{M}{\varepsilon} \left(\mathbf{dim}_P(\mathcal{H}) \log \frac{M}{\varepsilon} + \log \frac{1}{\delta}\right)\right), \quad (6)$$

in place of the sample complexity

$$m(\varepsilon, \delta) = O\left(\frac{M}{\varepsilon} \left(\log |\mathcal{H}| + \log \frac{1}{\delta}\right)\right)$$

derived from Theorem 1. Moreover, these bounds are distribution-independent, so \mathcal{P} can be taken to be the class of all probability distributions on Z .

We now give several examples to illustrate the use of this theorem. First, consider the standard PAC model in which the outcome space Y and the decision space A are both $\{0, 1\}$. In this case any loss function such that $l(0, 1) \neq l(0, 0)$ and $l(1, 1) \neq l(1, 0)$ is monotone over A , and in particular, the standard discrete loss function, $l(y, a) = 1$ if $y \neq a$, else $l(y, a) = 0$, is monotone. Clearly $M = 1$ in this case. Thus from Eq. (6) above we get a sample complexity bound of

$$O\left(\frac{1}{\varepsilon} \left(\mathbf{dim}_P(\mathcal{H}) \log \frac{1}{\varepsilon} + \log \frac{1}{\delta}\right)\right).$$

As mentioned in the previous section, $\mathbf{dim}_P(\mathcal{H})$ is the same as the Vapnik–Chervonenkis dimension of \mathbf{H} in this case; hence this bound is, up to constants, the same as that given in Theorem 2.1 of Blumer *et al.* (1989). A number of applications of this result are outlined in Blumer *et al.* (1989). Further applications, specifically for learning problems that have been studied recently in the mainstream artificial intelligence work, are given in Haussler (1988, 1989).

For our second example, consider the case where the outcome space Y is $\{0, 1\}$ but the decision space A is $[0, 1]$. Assume that the loss function is $l(y, a) = |a - y|^q$ for some $q > 0$. This case was examined with $q = 2$ by Kearns and Schapire (1990) in their investigation into the learnability of p -concepts. They showed that $\mathbf{dim}_P(l_{\mathcal{H}}) = \mathbf{dim}_P(\mathcal{H})$ in this case. Since the

loss l is monotone in A for all $q > 0$, Lemma 5 shows that this result holds for other values of q as well. Hence the conditions of Theorem 8 are met. Some applications are given in Kearns and Schapire (1990). (It should be noted that it is important that $A = [0, 1]$ in this case. The result does not hold in general for larger A .)

For our third example, consider the problem of logistic regression, as described in Section 1.1.3. In the simplest case the outcome space again has only two values, denoted y_1 and y_2 , where y_1 indicates that some event has taken place and y_2 indicates that it has not, and an action a represents an estimate of the log odds ratio $\ln(P(y_1)/P(y_2))$, where the probability P is conditioned on the observed instance x . Here $A = \mathfrak{R}$ and the log likelihood loss function is the logistic loss function, defined by $l(y_1, a) = \ln(1 + e^a) - a$ and $l(y_2, a) = \ln(1 + e^a)$. Again, it is easily verified that l is monotone in A . In logistic regression the standard assumption is that the instance space X is contained in \mathfrak{R}^n for some $n \geq 1$ and \mathcal{H} is contained in the family of all linear functions on X (see, e.g., McCullagh and Nelder (1989)). In this case, $\dim_p(\mathcal{H}) \leq n + 1$ by Theorem 4. By restricting A to a bounded range, we can then apply Theorem 8 to obtain sample complexity bounds that are linear in n .

As a last example, consider the problem of density estimation, as described in Section 1.1.4. Here there is only one outcome in Y , $A \subset \mathfrak{R}^+$, and $l(y, a) = l(a) = -\log a$. Thus clearly l is monotone in A . Thus we can apply Theorem 8 to the problem of density estimation as well, whenever the family of densities \mathcal{H} is uniformly bounded (away from zero) and has finite pseudo dimension.

6. CAPACITY AND METRIC DIMENSION OF FUNCTION CLASSES

In Sections 4 and 5 we showed how the pseudo dimension can be used to obtain distribution independent bounds on the random covering numbers needed for Theorem 2, thereby obtaining bounds on the sample size needed for uniform convergence and learning results. In this section we develop an alternate way of obtaining distribution independent bounds on random covering numbers. This method can sometimes be used in conjunction with the method given in the previous sections to extend that method to cover cases where the decision space A is not contained in \mathfrak{R} . We will demonstrate this in our analysis of the sample size needed for learning in feedforward neural networks in the following section.

The key idea is to introduce a pseudo metric (see Section 9.1) on the decision space A . The distance between two actions is the maximum difference in loss for these actions, over all possible outcomes.

DEFINITION 6. For every loss function $l: Y \times A \rightarrow [0, M]$, by ρ_l we denote the pseudo metric on A defined by $\rho_l(a, b) = \sup_{y \in Y} |l(y, a) - l(y, b)|$ for all $a, b \in A$.

Note that (A, ρ_l) is a bounded pseudo metric space: no two actions in A are more than M apart.

Since decision rules in \mathcal{H} map from the instance space X into A , the pseudo metric ρ_l on A can be used to induce a pseudo metric on \mathcal{H} in which two decision rules differ only to the extent that the actions that they prescribe differ with respect to ρ_l . There are several ways to do this. The easiest is to use an L^∞ function distance on \mathcal{H} , defining the distance between decision rules f and g as the supremum of $\rho_l(f(x), g(x))$ over all $x \in X$. This works, and is a useful method of obtaining uniform convergence and learning results (see related techniques used in White, 1990a). However, as we will see, the crucial issue is the size of the smallest ε -cover of the resulting pseudo metric space \mathcal{H} . In some cases we can get smaller covers, and hence better results, by using an L^1 function distance instead. Since the L^1 distance is never more than the L^∞ distance, the results are never worse. Thus we present this more powerful method here.

DEFINITION 7. Let \mathcal{H} be a family of functions from a set X into a bounded pseudo metric space (A, ρ) . Let P be a probability measure on X . Then $d_{L^1(P, \rho)}$ is the pseudo metric on \mathcal{H} defined by

$$d_{L^1(P, \rho)}(f, g) = \mathbf{E}(\rho(f(x), g(x))) = \int_X \rho(f(x), g(x)) dP(x)$$

for all $f, g \in \mathcal{H}$. For every $\varepsilon > 0$ let

$\mathcal{C}(\varepsilon, \mathcal{H}, \rho) = \sup\{\mathcal{N}(\varepsilon, \mathcal{H}, d_{L^1(P, \rho)})\}$ over all probability measures P on X .

If $\mathcal{N}(\varepsilon, \mathcal{H}, d_{L^1(P, \rho)})$ is infinite for some measure P , or if the set in this supremum is unbounded, then $\mathcal{C}(\varepsilon, \mathcal{H}, \rho) = \infty$. We call¹³ $\mathcal{C}(\varepsilon, \mathcal{H}, \rho)$ the *capacity* of \mathcal{H} . In analogy with the definition of metric dimension in Section 9.1, we define the *upper metric dimension* of \mathcal{H} by

$$\overline{\dim}(\mathcal{H}) = \limsup_{\varepsilon \rightarrow 0} \frac{\log \mathcal{C}(\varepsilon, \mathcal{H}, \rho)}{\log(1/\varepsilon)},$$

¹³ The term *metric entropy* is often used for the quantities $\log \mathcal{N}(\varepsilon, \mathcal{H}, d_{L^1(P, \rho)})$ and $\log \mathcal{C}(\varepsilon, \mathcal{H}, \rho)$ (Dudley, 1987; Quiroz, 1989). It is also used for an analogous, but fundamentally distinct, concept in the dynamical systems literature (e.g., Farmer, 1982). The term *capacity* has also been used with many other related meanings (Mandelbrot, 1982; Vapnik, 1982; Kolmogorov and Tihomirov, 1961; Farmer *et al.*, 1983; Baum and Haussler, 1989). Our usage here is taken from Dudley (1984).

and the *lower metric dimension*, denoted by $\underline{\dim}(\mathcal{H})$, is defined similarly using \liminf . When $\overline{\dim}(\mathcal{H}) = \underline{\dim}(\mathcal{H})$, then this quantity is denoted $\dim(\mathcal{H})$, and referred to simply as the *metric dimension* of \mathcal{H} . If $\mathcal{C}(\varepsilon, \mathcal{H}, \rho) = \infty$ for some $\varepsilon > 0$ then $\dim(\mathcal{H}) = \infty$.

We now show how bounds on the capacity of \mathcal{H} lead to distribution-independent bounds on the rate of uniform convergence of empirical risk estimates for functions in \mathcal{H} with respect to the loss function l . As before, let $Z = X \times Y$, let P be a probability distribution on Z , and let $l_{\mathcal{H}}$ be the family of functions on Z defined by $l_{\mathcal{H}} = \{l_h : h \in \mathcal{H}\}$, where $l_h(x, y) = l(y, h(x))$. Let $P_{|X}$ be the marginal on X of the joint distribution P on $X \times Y$ (see Section 1.5).

LEMMA 6. For all $\varepsilon > 0$,

$$\mathcal{N}(\varepsilon, l_{\mathcal{H}}, d_{L^1(P)}) \leq \mathcal{N}(\varepsilon, \mathcal{H}, d_{L^1(P_{|X}, \rho_l)}).$$

Proof. For every $h \in \mathcal{H}$ let $\psi(h) = l_h$. Hence ψ maps from \mathcal{H} onto $l_{\mathcal{H}}$. It suffices to show that ψ is a contraction, i.e., that

$$\forall f, g \in \mathcal{H}, \quad d_{L^1(P)}(\psi(f), \psi(g)) \leq d_{L^1(P_{|X}, \rho_l)}(f, g).$$

Let f and g be any two functions in \mathcal{H} . Then

$$\begin{aligned} d_{L^1(P)}(\psi(f), \psi(g)) &= \int_Z |l(y, f(x)) - l(y, g(x))| dP(x, y) \\ &\leq \int_Z \rho_l(f(x), g(x)) dP(x, y) \\ &= \int_X \rho_l(f(x), g(x)) dP_{|X}(x) \\ &= d_{L^1(P_{|X}, \rho_l)}(f, g). \quad \blacksquare \end{aligned}$$

This gives the following theorem about distribution-independent uniform convergence of risk estimates for learning.

THEOREM 9. Assume that the decision rule space \mathcal{H} and the loss function l are such that $l_{\mathcal{H}}$ is permissible. Let P be any probability distribution on $Z = X \times Y$. Assume that $m \geq 1$, $v > 0$, and $0 < \alpha < 1$. Let \vec{z} be generated by m independent draws from Z according to P . Then

$$\Pr(\exists h \in \mathcal{H} : d_v(\hat{\mathbf{r}}_{h,l}(\vec{z}), \mathbf{r}_{h,l}(P)) > \alpha) \leq 4\mathcal{C}(\alpha v/8, \mathcal{H}, \rho_l) e^{-\alpha^2 v m/8M}.$$

Proof. Let $\mathbf{F} = l_{\mathcal{H}}$. For any sequence \vec{z} of points in Z there is a trivial isometry between $(\mathbf{F}_{|\vec{z}}, d_{L^1})$ and $(\mathbf{F}, d_{L^1(P_{\vec{z}})})$, where $P_{\vec{z}}$ is the empirical

measure induced by \vec{z} , in which each set has measure equal to the fraction of the points in \vec{z} it contains. Thus by Lemma 6 above, we have

$$\mathcal{N}(\varepsilon, \mathbf{F}_{|\vec{z}}, d_{L^1}) = \mathcal{N}(\varepsilon, \mathbf{F}, d_{L^1(P_{\vec{z}})}) \leq \mathcal{N}(\varepsilon, \mathcal{H}, d_{L^1(P_{\vec{z}|X}, \rho_I)}) \leq \mathcal{C}(\varepsilon, \mathcal{H}, \rho_I)$$

for all $\vec{z} \in Z^{2m}$. Hence, setting $\varepsilon = \alpha v/8$, the given probability is at most $4\mathcal{C}(\alpha v/8, \mathcal{H}, \rho_I) e^{-\alpha^2 v m/8M}$ by Theorem 3. ■

In order to apply the above theorem, we need tools for bounding the capacity of various decision rule spaces. Along these lines, we close this section by proving two basic lemmas, one about the capacity of the free product of a set of function classes, and the other about the capacity of compositions of function classes.

DEFINITION 8. Let $(A_1, \rho_1), \dots, (A_k, \rho_k)$ be bounded metric spaces. Let $A = A_1 \times \dots \times A_k$ and let ρ be the metric on A defined by

$$\rho(\vec{u}, \vec{v}) = \frac{1}{k} \sum_{j=1}^k \rho_j(u_j, v_j)$$

for any $\vec{u} = (u_1, \dots, u_k)$ and $\vec{v} = (v_1, \dots, v_k) \in A$. For each j , $1 \leq j \leq k$, let \mathcal{H}_j be a family of functions from X into A_j . The *free product* of \mathcal{H}_1 through \mathcal{H}_k is the class of functions

$$\mathcal{H} = \{(f_1, \dots, f_k): f_j \in \mathcal{H}_j, 1 \leq j \leq k\},$$

where $(f_1, \dots, f_k): X \rightarrow A$ is the function defined by

$$(f_1, \dots, f_k)(x) = (f_1(x), \dots, f_k(x)).$$

LEMMA 7. If $\mathcal{H}, \mathcal{H}_1, \dots, \mathcal{H}_k$ are defined as above then

1. for any probability measure P on X and $\varepsilon > 0$,

$$\prod_{j=1}^k \mathcal{N}(2k\varepsilon, \mathcal{H}_j, d_{L^1(P, \rho_j)}) \leq \mathcal{N}(\varepsilon, \mathcal{H}, d_{L^1(P, \rho)}) \leq \prod_{j=1}^k \mathcal{N}(\varepsilon, \mathcal{H}_j, d_{L^1(P, \rho_j)});$$

2. $\overline{\dim}(\mathcal{H}) = \sum_{j=1}^k \overline{\dim}(\mathcal{H}_j)$, and similarly for $\underline{\dim}$ and \dim , when the latter is defined.

Proof. We begin with the second inequality of (1). For each $1 \leq j \leq k$ let U_j be an ε -cover for \mathcal{H}_j . Let

$$U = \prod_{j=1}^k U_j = \{(f_1, \dots, f_k): f_j \in U_j, 1 \leq j \leq k\}.$$

It suffices to show that U is an ε -cover for \mathcal{H} . Let $g = (g_1, \dots, g_k)$ be any function in \mathcal{H} . For each j , $1 \leq j \leq k$, find $f_j \in U_j$ such that $d_{L^1(P, \rho_j)}(f_j, g_j) \leq \varepsilon$. Let $f = (f_1, \dots, f_k)$. Then

$$\begin{aligned} d_{L^1(P, \rho)}(f, g) &= \int_X \frac{1}{k} \sum_{j=1}^k \rho_j(f_j(x), g_j(x)) dP(x) \\ &= \frac{1}{k} \sum_{j=1}^k \int_X \rho_j(f_j(x), g_j(x)) dP(x) \\ &= \frac{1}{k} \sum_{j=1}^k d_{L^1(P, \rho_j)}(f_j, g_j) \\ &\leq \varepsilon. \end{aligned}$$

Hence U is an ε -cover for \mathcal{H} .

The first inequality of (1) is verified similarly. For each $1 \leq j \leq k$ let V_j be a $k\varepsilon$ -separated subset of \mathcal{H}_j . Let $V = \prod_{j=1}^k V_j$. Let $f = (f_1, \dots, f_k)$ and $g = (g_1, \dots, g_k)$ be distinct functions in V . Then

$$d_{L^1(P, \rho)}(f, g) = \frac{1}{k} \sum_{j=1}^k d_{L^1(P, \rho_j)}(f_j, g_j) > \varepsilon.$$

Hence V is an ε -separated subset of \mathcal{H} . It follows that

$$\prod_{j=1}^k \mathcal{M}(k\varepsilon, \mathcal{H}_j, d_{L^1(P, \rho_j)}) \leq \mathcal{M}(\varepsilon, \mathcal{H}, d_{L^1(P, \rho)}).$$

The first inequality of (1) then follows using Theorem 12.

From (1) we have

$$\prod_{j=1}^k \mathcal{C}(2k\varepsilon, \mathcal{H}_j, \rho_j) \leq \mathcal{C}(\varepsilon, \mathcal{H}, \rho) \leq \prod_{j=1}^k \mathcal{C}(\varepsilon, \mathcal{H}_j, \rho_j).$$

Part (2) follows easily from this. ■

DEFINITION 9. Let P be a probability measure on X and let f be a measurable function from X into Y . Then P_f denotes the probability measure on Y induced by f ; i.e.,

$$P_f(S) = P(f^{-1}(S)) \quad \text{for all measurable } S \subset Y.$$

DEFINITION 10. Let f be a function from a metric space (X, ρ) into a metric space (Y, σ) . A *Lipschitz bound* on f is a real number $b > 0$ such that for all $x, y \in X$, $\sigma(f(x), f(y)) \leq b\rho(x, y)$. The Lipschitz bound on f is the smallest such b . If \mathcal{F} is a class of functions from (X, ρ) into (Y, σ) then b is a *uniform Lipschitz bound* on \mathcal{F} if b is Lipschitz bound on f for all $f \in \mathcal{F}$.

LEMMA 8. Let $(X_1, \rho_1), \dots, (X_{k+1}, \rho_{k+1})$ be metric spaces, where (X_j, ρ_j) is bounded, $2 \leq j \leq k$, and \mathcal{H}_j be a class of functions with $f: X_j \rightarrow X_{j+1}$ for all $f \in \mathcal{H}_j$, $1 \leq j \leq k$. Let b_j be a uniform Lipschitz bound on \mathcal{H}_j for all $2 \leq j \leq k$. Let \mathcal{H} denote the class of all functions from X_1 into X_{k+1} defined by compositions of functions in the \mathcal{H}_j 's; i.e.,

$$\mathcal{H} = \{f_k \circ f_{k-1} \circ \dots \circ f_1 : f_j \in \mathcal{H}_j, 1 \leq j \leq k\}.$$

1. For any $\varepsilon, \varepsilon_1, \dots, \varepsilon_k > 0$ such that

$$\varepsilon = \sum_{j=1}^k \left(\prod_{l=j+1}^k b_l \right) \varepsilon_j$$

we have

$$\mathcal{C}(\varepsilon, \mathcal{H}, \rho_{k+1}) \leq \prod_{j=1}^k \mathcal{C}(\varepsilon_j, \mathcal{H}_j, \rho_{j+1}).$$

2. $\overline{\dim}(\mathcal{H}) \leq \sum_{j=1}^k \overline{\dim}(\mathcal{H}_j)$, and similarly for $\underline{\dim}$ and \dim , when the latter is defined.

Proof. Fix a probability measure P on X_1 . We define a tree-structured family of covers for the \mathcal{H}_j 's by induction as follows. For the basis case, let U be a minimum-sized ε_1 -cover for \mathcal{H}_1 w.r.t. the measure P on X_1 ; i.e., $|U| = \mathcal{N}(\varepsilon_1, \mathcal{H}_1, d_{L^1(P, \rho_2)})$ and every function in \mathcal{H}_1 is $L^1(P, \rho_2)$ -approximated to within ε_1 by some function in U . Now for each j , $2 \leq j \leq k$, and for each sequence of functions f_1, \dots, f_{j-1} , where $f_1 \in U$, $f_2 \in U_{f_1}$, $f_3 \in U_{f_1, f_2}$, ..., $f_{j-1} \in U_{f_1, \dots, f_{j-2}}$, let $U_{f_1, \dots, f_{j-1}}$ be a minimum-sized ε_j cover for \mathcal{H}_j w.r.t. the L^1 metric for the measure $P_{f_{j-1} \circ f_{j-2} \circ \dots \circ f_1}$ on X_j and the metric ρ_{j+1} on X_{j+1} .

Next we define a cover V for \mathcal{H} by composing functions in the covers for the \mathcal{H}_j 's. If $k = 1$ then $V = U$. Otherwise

$$V = \{f_k \circ f_{k-1} \circ \dots \circ f_1 : f_1 \in U, f_2 \in U_{f_1}, f_3 \in U_{f_1, f_2}, \dots, \text{ and } f_k \in U_{f_1, \dots, f_{k-1}}\}.$$

Since $\mathcal{N}(\varepsilon_j, \mathcal{H}_j, d_{L^1(P_{f_{j-1} \circ \dots \circ f_1}, \rho_{j+1})}) \leq \mathcal{C}(\varepsilon_j, \mathcal{H}_j, \rho_{j+1})$ for all $1 \leq j \leq k$ and all f_1, \dots, f_{j-1} , it is clear that $|V| \leq \prod_{j=1}^k \mathcal{C}(\varepsilon_j, \mathcal{H}_j, \rho_{j+1})$. Hence it remains to show that V is an ε -cover for \mathcal{H} .

Suppose that $g = g_k \circ g_{k-1} \circ \dots \circ g_1 \in \mathcal{H}$. Find

- $f_1 \in U$ such that $d_{L^1(P, \rho_2)}(f_1, g_1) \leq \varepsilon_1$,
- $f_2 \in U_{f_1}$ such that $d_{L^1(P_{f_1}, \rho_3)}(f_2, g_2) \leq \varepsilon_2$, ..., and
- $f_k \in U_{f_1, \dots, f_{k-1}}$ such that $d_{L^1(P_{f_{k-1} \circ \dots \circ f_1}, \rho_{k+1})}(f_k, g_k) \leq \varepsilon_k$.

Let $f = f_k \circ f_{k-1} \circ \cdots \circ f_1 \in V$. It suffices to show that $d_{L^1(P, \rho_{k+1})}(f, g) \leq \varepsilon$. We prove that for all h , $1 \leq h \leq k$,

$$d_{L^1(P, \rho_{h+1})}(f_h \circ \cdots \circ f_1, g_h \circ \cdots \circ g_1) \leq \sum_{j=1}^h \left(\prod_{l=j+1}^h b_l \right) \varepsilon_j.$$

Since

$$\varepsilon = \sum_{j=1}^k \left(\prod_{l=j+1}^k b_l \right) \varepsilon_j,$$

part (1) of the result follows.

If $h = 1$ then the result follows directly from our definition of f . Otherwise

$$\begin{aligned} & d_{L^1(P, \rho_{h+1})}(f_h \circ \cdots \circ f_1, g_h \circ \cdots \circ g_1) \\ &= \int_{X_1} \rho_{h+1}(f_h \circ \cdots \circ f_1(x), g_h \circ \cdots \circ g_1(x)) dP(x) \\ &\leq \int_{X_1} \rho_{h+1}(g_h \circ f_{h-1} \circ \cdots \circ f_1(x), g_h \circ g_{h-1} \circ \cdots \circ g_1(x)) dP(x) \\ &\quad + \int_{X_1} \rho_{h+1}(f_h \circ f_{h-1} \circ \cdots \circ f_1(x), g_h \circ f_{h-1} \circ \cdots \circ f_1(x)) dP(x) \\ &\leq b_h \int_{X_1} \rho_h(f_{h-1} \circ \cdots \circ f_1(x), g_{h-1} \circ \cdots \circ g_1(x)) dP(x) \\ &\quad + \int_{X_h} \rho_{h+1}(f_h(y), g_h(y)) dP_{f_{h-1}, \dots, f_1}(y) \text{ (by the Lipschitz assumption)} \\ &\leq b_h \sum_{j=1}^{h-1} \left(\sum_{l=j+1}^{h-1} b_l \right) \varepsilon_j + \varepsilon_h \text{ (by the inductive hypothesis and the definition of } f_h) \\ &= \sum_{j=1}^h \left(\prod_{l=j+1}^h b_l \right) \varepsilon_j. \end{aligned}$$

To prove part (2), let $a_j = \prod_{l=j+1}^k b_l$ and set $\varepsilon_j = \varepsilon/ka_j$ for $1 \leq j \leq k$. By part (1), $\mathcal{C}(\varepsilon, \mathcal{H}, \rho_{k+1}) \leq \prod_{j=1}^k \mathcal{C}(\varepsilon_j, \mathcal{H}_j, \rho_{j+1})$. Thus

$$\begin{aligned} \overline{\dim}(\mathcal{H}) &= \limsup_{\varepsilon \rightarrow 0} \left(\frac{\log(\mathcal{C}(\varepsilon, \mathcal{H}, \rho_{k+1}))}{\log(1/\varepsilon)} \right) \\ &\leq \limsup_{\varepsilon \rightarrow 0} \left(\frac{\log(\prod_{j=1}^k \mathcal{C}(\varepsilon/ka_j, \mathcal{H}_j, \rho_{j+1}))}{\log(1/\varepsilon)} \right) \\ &= \sum_{j=1}^k \limsup_{\varepsilon \rightarrow 0} \left(\frac{\log(\mathcal{C}(\varepsilon/ka_j, \mathcal{H}_j, \rho_{j+1}))}{\log(1/\varepsilon)} \right) \end{aligned}$$

$$\begin{aligned}
&= \sum_{j=1}^k \limsup_{\varepsilon_j \rightarrow 0} \left(\frac{\log(\mathcal{C}(\varepsilon_j, \mathcal{H}_j, \rho_{j+1}))}{\log(1/\varepsilon_j) + \log(1/ka_j)} \right) \\
&= \sum_{j=1}^k \limsup_{\varepsilon_j \rightarrow 0} \left(\frac{\log(\mathcal{C}(\varepsilon_j, \mathcal{H}_j, \rho_{j+1}))}{\log(1/\varepsilon_j)} \right) \\
&= \sum_{j=1}^k \overline{\dim}(\mathcal{H}_j). \quad \blacksquare
\end{aligned}$$

7. SAMPLE SIZE BOUNDS FOR LEARNING WITH MULTI-LAYER NEURAL NETS

We now present some applications of the results of the previous section to learning with feedforward neural nets (see, e.g., Rumelhart and McClelland, 1986; Poggio and Girosi, 1989). The decision rule space \mathcal{H} represented by a feedforward neural net consists of a family of functions from an instance space $X \subset \mathfrak{R}^n$ into a decision space $A \subset \mathfrak{R}^k$ for some k , $n \geq 1$. To apply Theorem 9 of the previous section, we need to obtain an upper bound on the capacity $\mathcal{C}(\varepsilon, \mathcal{H}, \rho_l)$ of such decision rule spaces for various loss functions l .

For many loss functions, the metric ρ_l on $A \subset \mathfrak{R}^k$ can be bounded in terms of the d_{L^1} metric; i.e., we can find a constant c_l such that for all $\vec{a} = (a_1, \dots, a_k)$ and $\vec{b} = (b_1, \dots, b_k)$ in A , $\rho_l(\vec{a}, \vec{b}) \leq c_l d_{L^1}(\vec{a}, \vec{b}) = (c_l/k) \sum_{i=1}^k |a_i - b_i|$. In this case it is clear that $\mathcal{C}(\varepsilon, \mathcal{H}, \rho_l) \leq \mathcal{C}(\varepsilon/c_l, \mathcal{H}, d_{L^1})$. Thus our problem is reduced to obtaining an upper bound on the capacity $\mathcal{C}(\varepsilon/c_l, \mathcal{H}, d_{L^1})$.

We now give a few examples to illustrate this reduction. First consider the common case in which the outcome space Y is also contained in \mathfrak{R}^k ; e.g., we receive explicit feedback on each coordinate of our action $\vec{a} \in A \subset \mathfrak{R}^k$. This occurs when each coordinate a_i of the action \vec{a} is a prediction of the corresponding coordinate of the outcome \vec{y} . Here the loss function l may itself be a metric on \mathfrak{R}^k which measures the distance between the predicted vector and the actual outcome vector. When l is a metric, we have for any actions $\vec{a}, \vec{b} \in A$

$$\rho_l(\vec{a}, \vec{b}) = \sup_{\vec{y} \in Y} |l(\vec{y}, \vec{a}) - l(\vec{y}, \vec{b})| \leq l(\vec{a}, \vec{b})$$

by the triangle inequality for l . Thus if the metric l is bounded with respect to the d_{L^1} metric, i.e., $l(\vec{a}, \vec{b}) \leq c_l d_{L^1}(\vec{a}, \vec{b})$ for all $\vec{a}, \vec{b} \in A$, then we have $\rho_l(\vec{a}, \vec{b}) \leq c_l d_{L^1}(\vec{a}, \vec{b})$. For example, if $l(\vec{y}, \vec{a}) = d_{L^2}(\vec{y}, \vec{a}) = ((1/k) \sum_{i=1}^k (y_i - a_i)^2)^{1/2}$ then we may take $c_l = 1$, and similarly for the other d_{L^q} metrics, for $q > 1$.

Note that the above trick does not apply to the mean squared loss $l(\vec{y}, \vec{a}) = (1/k) \sum_{i=1}^k (y_i - a_i)^2$ since this loss does not satisfy the triangle

inequality. However, in this case it is easy to show by direct calculation that if the outcome space Y is bounded, e.g., $Y \subset [0, M]^k$, then $\rho_l(\vec{a}, \vec{b}) \leq 2M d_{L^1}(\vec{a}, \vec{b})$, and hence we may take $c_l = 2M$.

For our final example, consider the case when $Y = \{0, 1\}^k$, $A \subset [0, 1]^k$, and l is the cross entropy loss,

$$l(\vec{y}, \vec{a}) = - \sum_{i=1}^k (y_i \ln a_i + (1 - y_i) \ln(1 - a_i)).$$

As discussed in Section 1.1.3, this is the log likelihood loss for the regression problem in which the action \vec{a} represents a vector of probabilities for independent Bernoulli variables, and the outcome \vec{y} gives the observed values of these variables. This loss is bounded if we restrict the probabilities in \vec{a} to be between B and $1 - B$ for some $0 < B \leq \frac{1}{2}$. In this case

$$\begin{aligned} \rho_l(\vec{a}, \vec{b}) &= \sup_{\vec{y} \in Y} \left| \sum_{i=1}^k \left(y_i \ln \frac{b_i}{a_i} + (1 - y_i) \ln \frac{(1 - b_i)}{(1 - a_i)} \right) \right| \\ &\leq \sum_{i=1}^k \left| \ln \frac{b_i}{a_i} \right| + \sum_{i=1}^k \left| \ln \frac{(1 - b_i)}{(1 - a_i)} \right| \\ &\leq \frac{2}{B} \sum_{i=1}^k |a_i - b_i|. \end{aligned}$$

The latter inequality follows from the fact that for $x, y > 0$,

$$\left| \ln \frac{x}{y} \right| = \ln \frac{\max(x, y)}{\min(x, y)} \leq \frac{\max(x, y)}{\min(x, y)} - 1 = \frac{|x - y|}{\min(x, y)}.$$

Thus in this case we may take $c_l = 2k/B$.

We now turn to the task of obtaining an upper bound on the capacity $\mathcal{C}(\varepsilon, \mathcal{H}, d_{L^1})$ when the decision rules in \mathcal{H} map into a decision space $A \subset \mathbb{R}^k$, and in particular, when these decision rules are represented by neural networks. When $k = 1$, i.e., the neural net has only one output, the decision rule space \mathcal{H} is a family of real-valued functions and $d_{L^1}(a, b) = |a - b|$ for $a, b \in A$. In this case we can apply the methods and results of Section 4. We must first find an upper bound on $\dim_{\mathbf{P}}(\mathcal{H})$, the pseudo dimension of \mathcal{H} . Then, when A is bounded, from the bound on $\dim_{\mathbf{P}}(\mathcal{H})$ we get a bound on the capacity $\mathcal{C}(\varepsilon, \mathcal{H}, d_{L^1})$ using Theorem 6. This also gives a bound on the metric dimension of \mathcal{H} .

THEOREM 10. *Let \mathcal{H} be a family of functions from X into $A = [0, M]$. Assume $\dim_{\mathbf{P}}(\mathcal{H}) = d$ for some $1 \leq d < \infty$.*

1. For all $0 < \varepsilon \leq M$,

$$\mathcal{C}(\varepsilon, \mathcal{H}, d_{L^1}) < 2 \left(\frac{2eM}{\varepsilon} \ln \frac{2eM}{\varepsilon} \right)^d.$$

2. $\overline{\dim}(\mathcal{H}) = \dim_P(\mathcal{H})$.

Proof. Let P be any probability measure on X . Then by Theorems 12 and 6 in Sections 9.1 and 4,

$$\mathcal{N}(\varepsilon, \mathcal{H}, d_{L^1(P)}) \leq \mathcal{M}(\varepsilon, \mathcal{H}, d_{L^1(P)}) < 2 \left(\frac{2eM}{\varepsilon} \ln \frac{2eM}{\varepsilon} \right)^d.$$

(Theorem 6 is applied with $Z = X$ and $F = \mathcal{H}$.) This gives (1), and (2) follows easily from (1). ■

In the general case, where $A \subset \mathfrak{R}^k$ for $k > 1$, we can apply the methods from the previous section, in addition to the pseudo dimension methods from Section 4, to obtain bounds on $\mathcal{C}(\varepsilon, \mathcal{H}, d_{L^1})$. We illustrate this for the case where \mathcal{H} is the class of decision rules represented by a feedforward neural network.

A feedforward neural network is defined as a directed acyclic graph in which the incoming edges to each node (or *unit*) are ordered and each incoming edge can carry a real number representing the *activation* on that edge. We will assume that all activations are restricted to the interval $[c_0, c_1]$ for some constants $c_0 < c_1$. The units are divided into *input units*, which have no incoming edges from other units and serve as input ports for the network (their activations are determined by these external inputs), and *computation units*, which have incoming edges from other units and compute an activation based on the activations on these incoming edges. After an activation has been determined, this activation is placed on the outgoing edges of the unit. Computation units with no outgoing edges are called *output units* and serve as output ports for the network. Computation units that are not output units are called *hidden units*. The network as a whole computes a function that maps from vectors of activation values in its input units to vectors of activation values in its output units by composing the functions computed by its computation units in the obvious way.

The action of a computation unit with n incoming edges can be specified by a function f from $[c_0, c_1]^n$ into $[c_0, c_1]$, where $f(\vec{x})$ is the resulting activation of the unit when the activations of its incoming edges are given by the vector $\vec{x} \in [c_0, c_1]^n$. In the nets we consider, the function f is defined by

$$f(\vec{x}) = \sigma \left(\mu(\phi_1(\vec{x}), \dots, \phi_k(\vec{x})) + \theta + \sum_{j=1}^k w_j \phi_j(\vec{x}) \right),$$

where the w_j 's are adjustable real *weights*, θ is an adjustable real *bias*, ϕ_1, \dots, ϕ_k are fixed real-valued functions which we call the *input transformers*, $\mu: \mathfrak{R}^k \rightarrow \mathfrak{R}$ is a fixed function which we call the *global modifier*, and $\sigma: \mathfrak{R} \rightarrow [c_0, c_1]$ is a fixed non-increasing or non-decreasing function which we call the *squashing function*. Different units can have different modifiers, transformers, and squashing functions. We say that the function f computed by a given computation unit with n incoming edges has *Lipschitz bound* b if for any $\vec{x}, \vec{y} \in [c_0, c_1]^n$, $|f(\vec{x}) - f(\vec{y})| \leq b d_L(\vec{x}, \vec{y})$.

We give a few examples to illustrate the flexibility of this model at the level of the individual computation unit. First assume that $k = n$, i.e., the number of input transformers is the same as the number of inputs, and that each input transformer simply extracts a component of the input, i.e., $\phi_j(\vec{x}) = x_j$, $1 \leq j \leq n$. In this case, which is the standard case for most neural net research, the overall input transformation is just the identity map and can be ignored. In this standard case, if the global modifier $\mu = 0$ we get what is known as a quasi-linear unit (Rumelhart and McClelland, 1986):

$$f(\vec{x}) = \sigma \left(\theta + \sum_{j=1}^k w_j x_j \right).$$

In the standard case, if $\mu(\vec{x}) = \sum_{j=1}^n x_j^2$ we get a unit that computes a function of the form

$$f(\vec{x}) = \sigma \left(\theta' + \sum_{j=1}^n (x_j - a_j)^2 \right),$$

where $a_j = -w_j/2$ and $\theta' = \theta - \sum_{j=1}^n a_j^2$. This is similar to what is called a *radial basis* unit in the neural net literature (Poggio and Girosi, 1989; Moody and Darken, 1989).

Now assume that $k = n$ but the input transformers take logs of the components of the inputs; i.e., $\phi_j(\vec{x}) = \log x_j$. (Here we assume $c_0 > 0$.) Let $\mu = 0$ and change the squashing function σ to σ' , where $\sigma'(x) = \sigma(e^x)$. Then

$$f(\vec{x}) = \sigma' \left(\theta + \sum_{j=1}^n w_j \log x_j \right) = \sigma \left(e^\theta \prod_{j=1}^n x_j^{w_j} \right),$$

giving what is commonly known as a *product unit* (Durbin and Rumelhart, 1989).

We define a feedforward *architecture* as a feedforward net with unspecified weights and biases; i.e., each computation unit has a fixed global modifier, a fixed squashing function and fixed input transformers, but it has variable weights and a variable bias. We say a unit is *at depth*

j in an architecture if the longest (directed) path from an input unit to that unit has j edges. Thus all input units are at depth 0, all computation units that have incoming edges only from input units are at depth 1, all computation units that have incoming edges only from input units and computation units at depth 1 are at depth 2, etc. The *depth of the architecture* is the depth of the deepest unit in it.

We can bound the capacity of the decision rule space represented by a feedforward architecture as follows.

THEOREM 11. *Let \mathcal{A} be a feedforward architecture as above with $n \geq 1$ input units, $k \geq 1$ output units, and depth $d \geq 1$. Let W be the total number of adjustable weights and biases in \mathcal{A} . Assume $b_j \geq 1$ for $2 \leq j \leq d$, and let \mathcal{H} be all functions from $[c_0, c_1]^n$ into $[c_0, c_1]^k$ representable on \mathcal{A} by setting the adjustable weights and biases such that for all j , $2 \leq j \leq d$, the average of the Lipschitz bounds of the functions computed by computation units at depth j is at most b_j . Then for all $0 < \varepsilon \leq c_2 - c_1$,*

$$\mathcal{C}(\varepsilon, \mathcal{H}, d_{L^1}) \leq \left(\frac{2e(c_2 - c_1) d \prod_{l=2}^d b_l}{\varepsilon} \right)^{2W}. \quad \blacksquare$$

Proof. For each j , $0 \leq j \leq d$, let n_j be the number of units at depth j in the architecture \mathcal{A} . For each j , $0 \leq j \leq d-1$, let $l_j = \sum_{i=0}^j n_j$, and let $l_d = n_d = k$. For each j , $1 \leq j \leq d+1$, let $X_j = [c_1, c_2]^{l_{j-1}}$. Then for each j , $1 \leq j \leq d$, we can define the family \mathcal{H}_j of functions from X_j into X_{j+1} in the following manner.

First assume $j < d$. Let u_1, \dots, u_{n_j} be an enumeration of the computation units at depth j and f_1, \dots, f_{n_j} be functions such that f_i can be represented by u_i , $1 \leq i \leq n_j$, and the average Lipschitz bound on the f_i s is at most b_j . Let h_j be the free product of f_1, \dots, f_{n_j} and l_{j-1} copies of the identity function on $[c_1, c_2]$. Thus $h_j: X_j \rightarrow X_{j+1}$. The function h_j represents a mapping from the sequence of all activations of units at depth at most $j-1$ to the sequence of all activations of units at depth at most j , where the activations at depth at most $j-1$ are unaltered, and the new activations, i.e., those at depth j , are calculated by f_1, \dots, f_{n_j} . The family \mathcal{H}_j consists of all functions h_j obtained in this manner, by varying the weights and biases in the units u_1, \dots, u_{n_j} at depth j in such a manner that the Lipschitz constraint is satisfied.

When $j = d$, no subsequent calculations will be performed so we no longer need to preserve the activations of shallower units. Hence, we omit the identity function components in each $h_d \in \mathcal{H}_d$. Otherwise the definition of \mathcal{H}_d is the same as that for \mathcal{H}_j , where $j < d$.

It is clear that the class \mathcal{H} in the statement of the theorem can be represented as the class of compositions of functions from classes $\mathcal{H}_1, \dots, \mathcal{H}_d$. Since the identity function has Lipschitz bound $1 \leq b_j$, the

average Lipschitz bound on the components of each function $h_j \in \mathcal{H}_j$ is at most b_j . It is easily verified that a free product function is Lipschitz bounded by the average of the Lipschitz bounds on its component functions. Hence by assumption, b_j is a uniform Lipschitz bound on \mathcal{H}_j , $2 \leq j \leq d$. For each j , $1 \leq j \leq d$, let $a_j = \prod_{l=j+1}^d b_l$ and $\varepsilon_j = \varepsilon / d a_j$. Since $\varepsilon < c_2 - c_1$ and $a_j \geq 1$, $\varepsilon_j \leq c_2 - c_1$. Let ρ_j be the d_{L^1} metric on X_j , $1 \leq j \leq d+1$. Then by Lemma 8, part (1),

$$\mathcal{C}(\varepsilon, \mathcal{H}, d_{L^1}) \leq \prod_{j=1}^d \mathcal{C}(\varepsilon_j, \mathcal{H}_j, \rho_{j+1}).$$

For each j , \mathcal{H}_j is contained in the free product of l_j function classes. Each class \mathcal{F} in this product is either the trivial class containing only the identity function, or is a finite dimensional vector space of real-valued functions, summed with a fixed modifier and then composed with a non-increasing or non-decreasing squashing function. In the latter case, the dimension N of this vector space is the number of free parameters associated with the corresponding computation unit, i.e., the number of weights plus one (for the adjustable bias). Hence by Theorems 4 and 5 in Section 4, the pseudo dimension $\mathbf{dim}_p(\mathcal{F}) \leq N$. Thus by Theorem 10 above,

$$\mathcal{C}(\varepsilon_j, \mathcal{F}, d_{L^1}) \leq 2 \left(\frac{2e(c_2 - c_1)}{\varepsilon_j} \ln \frac{2e(c_2 - c_1)}{\varepsilon_j} \right)^N \leq \left(\frac{2e(c_2 - c_1)}{\varepsilon_j} \right)^{2N},$$

since $2 \ln x < x$ and $N \geq 1$. Since the capacity of a class with only one function is 1, it follows from Lemma 7 part (1) that

$$\mathcal{C}(\varepsilon_j, \mathcal{H}_j, \rho_{j+1}) \leq \left(\frac{2e(c_2 - c_1)}{\varepsilon_j} \right)^{2W_j},$$

where W_j is the total number of weights and biases of all computation nodes at depth j . Multiplying these bounds over all j , it follows that

$$\begin{aligned} \mathcal{C}(\varepsilon, \mathcal{H}, d_{L^1}) &\leq \prod_{j=1}^d \left(\frac{2e(c_2 - c_1)}{\varepsilon_j} \right)^{2W_j} \\ &= \prod_{j=1}^d \left(\frac{2e(c_2 - c_1) d \prod_{l=j+1}^d b_l}{\varepsilon} \right)^{2W_j} \\ &\leq \left(\frac{2e(c_2 - c_1) d \prod_{l=2}^d b_l}{\varepsilon} \right)^{2W}. \quad \blacksquare \end{aligned}$$

COROLLARY 3. *Let n, k, \mathcal{H}, W, d , and b_2, \dots, b_d be as in the previous theorem. Let X be the instance space $[c_1, c_2]^n$, let A be the decision space*

$[c_1, c_2]^k$, and let Y be any outcome space. Let $l: Y \times A \rightarrow [0, M]$ be a loss function and let c_l be a constant such that $\rho_l(\vec{a}, \vec{b}) \leq c_l d_{L^1}(\vec{a}, \vec{b})$ for all $\vec{a}, \vec{b} \in A$. Let $m \geq 1$, $0 < v \leq 8(c_2 - c_1)$, $0 < \alpha < 1$, and let P be any probability distribution on $Z = X \times Y$. Let \vec{z} be generated by m independent random draws from Z according to P .

1. Then

$$\begin{aligned} \Pr(\exists f \in \mathcal{H}: d_v(\hat{\mathbf{r}}_{f,l}(\vec{z}), \mathbf{r}_{f,l}(P)) > \alpha) \\ \leq 4 \left(\frac{c_l 16e(c_2 - c_1) d \prod_{l=2}^d b_l}{\alpha v} \right)^{2W} e^{-x^2 vm/8M}. \end{aligned}$$

2. Assume that for each computation unit at depth 2 and above the number of weights is at most W_{\max} , no weight is allowed to have absolute value greater than β , the input transformers are identity functions, the global modifier has Lipschitz bound at most r , and the squashing function has Lipschitz bound at most s , where $s(\beta W_{\max} + r) \geq 1$. Then for any $0 < \delta < 1$, the probability in (1) is less than δ for sample size

$$m = O \left(\frac{M}{\alpha^2 v} \left(W \left(\log \frac{c_l(c_2 - c_1)}{\alpha v} + d \log(s(\beta W_{\max} + r)) \right) + \log \frac{1}{\delta} \right) \right).$$

Proof. Let $\varepsilon = \alpha v/8 \leq c_2 - c_1$. It can be verified that $l_{\mathcal{H}}$ is permissible for the decision rule space \mathcal{H} . Hence, using Theorem 9 and Theorem 11,

$$\begin{aligned} \Pr(\exists f \in \mathcal{H}: d_v(\hat{\mathbf{r}}_{f,l}(\vec{z}), \mathbf{r}_{f,l}(P)) > \alpha) \\ \leq 4\mathcal{C}(\alpha v/8, \mathcal{H}, \rho_l) e^{-x^2 vm/8M} \\ \leq 4\mathcal{C}(\alpha v/8c_l, \mathcal{H}, d_{L^1}) e^{-x^2 vm/8M} \\ \leq 4 \left(\frac{c_l 16e(c_2 - c_1) d \prod_{l=2}^d b_l}{\alpha v} \right)^{2W} e^{-x^2 vm/8M}. \end{aligned}$$

For the second bound, it is readily verified that the L^1 Lipschitz bound for a linear function defined by W_{\max} weights and a bias is no more than W_{\max} times the largest absolute value of any weight. Furthermore, the Lipschitz bound for the sum of two functions is no more than the sum of the Lipschitz bounds on the individual functions, and the Lipschitz bound for the composition of two functions is no more than the product of the Lipschitz bounds on the individual functions. Thus if the input transformers are identity functions, the global modifier and squashing function have Lipschitz bounds r and s respectively and no weight is allowed to

have absolute value greater than β , then the Lipschitz bound for a computation unit is at most $s(\beta W_{\max} + r)$. If this holds for all units at depth 2 and above, may take $b_j = s(\beta W_{\max} + r)$ for all $j \geq 2$ in the first bound. Solving for m , this gives the order-of-magnitude estimate of the second bound. ■

We give the constants in the upper bound of part (1) of the above theorem only to show that they are not outlandishly large. We do not mean to suggest that the bound is tight. At present we cannot even verify that the asymptotic bound of part (2) is tight. In particular, we cannot show that the dependence on the Lipschitz bounds is necessary. Evidence that it may not be necessary comes from the analysis of the case where the squashing function σ is a sharp threshold function, i.e., $\sigma(x) = \text{sign}(x)$. Corollary 3 does not apply in this case, because the jump in σ prevents us from obtaining a Lipschitz bound on the computation units. As we let a smooth σ approach the sign function, its slope increases without limit, and thus the bound given in Corollary 3 degenerates. Nevertheless, using the techniques in Baum and Haussler (1989) it can be shown that results similar to Corollary 3 hold in this case, except that no Lipschitz bounds are required, and a bound on the sample size is

$$O\left(\frac{1}{\alpha^2 v} \left(W \log \frac{N}{\alpha v} + \log \frac{1}{\delta} \right)\right),$$

where N is the total number of computation units in the net. Details are given in Theorem 13 in the Appendix.

Despite the uncertainty about the need for the Lipschitz bounds, the result does give some indication of the maximum training sample size that will be needed for many popular network configurations. For example, if the squashing function is chosen as $\sigma(x) = 1/(1 + e^{-x/T})$ for some temperature $T > 0$ then it can be shown that the Lipschitz bound s for σ is $1/4T$. When the modifier $\mu \equiv 0$, then $r = 0$. Thus in this case the term $d \log(s(\beta W_{\max} + r))$ in the bound of Corollary 3 becomes $d \log(\beta W_{\max}/T)$. If the maximum weight β , the temperature T , and the depth d are constants, along with $c_2 - c_1$, M , and c_l , then the asymptotic bound of the theorem becomes

$$O\left(\frac{1}{\alpha^2 v} \left(W \log \frac{W_{\max}}{\alpha v} + \log \frac{1}{\delta} \right)\right),$$

which is similar to the bound obtained in Baum and Haussler (1989).

It should also be noted that Corollary 3 does have the feature that no Lipschitz bounds are required on the computation units at depth one. Thus

if all computation units are at depth one, i.e., there are no hidden units, then no Lipschitz units are required at all. If the architecture has only one layer of hidden units at depth one and a single output unit at depth two, as is quite common, then Lipschitz bounds are required only on the output unit. This means that the weights and biases associated with the hidden units do not need to be bounded in order to get the rates of uniform convergence given by Corollary 3, as they would, for example, if the methods given in White (1990a) were used to obtain a result of this type.

For an example of the above, consider networks that implement generalized radial basis¹⁴ functions, as described in Poggio and Girosi (1989). These networks have one layer of hidden units at depth 1 and one output unit at depth 2. The structure of the hidden units is as described in the example above: the input transformers are identity functions, the modifier is $\sum_{i=1}^n x_i^2$, and the squashing function is usually a smooth decreasing function. The output unit simply computes a weighted sum, so for this unit the modifier is the 0 function and the squashing function is the identity. Since this is the only unit at depth 2 and above, we require a Lipschitz bound only for this unit. If β is a bound on the maximum weight coming into the output unit, and W_{\max} is the number of units in the hidden layer, then the term $d \log(s(\beta W_{\max} + r))$ in the above bound becomes $\log(\beta W_{\max})$. Again, fixing β , $c_2 - c_1$, M , and c_l gives the same sample size bound,

$$O\left(\frac{1}{\alpha^2 v} \left(W \log \frac{W_{\max}}{\alpha v} + \log \frac{1}{\delta} \right)\right),$$

similar to that obtained in Baum and Haussler (1989).

Since W appears to be the dominant factor in these bounds, apart from the accuracy parameters α and v , these bounds support the conventional wisdom that the training set size should be primarily related to the number of adjustable parameters in the net. They also support the notion that this relationship between appropriate training size and the number of parameters is nearly linear, at least in the worst case. Further work is needed to sharpen these relationships (see, e.g., the lower bounds obtained in Baum and Haussler, 1989)).

¹⁴ The computation units in the network of radial basis functions described here are quite primitive in that they have no adjustable multiplicative parameter included in their basic radial distance calculation. Such parameters would be needed to do any reasonable type of kernel-based density estimation (see, e.g., Duda and Hart (1973). These parameters can be simulated by inserting another layer of computation units between the inputs and the layer described here. Alternately, the analysis can also be done directly for adjustable kernel units. This cleaner approach is detailed in Pollard (1986).

8. CONCLUSION

We have extended the PAC learning model to a more general decision theoretic framework so that it addresses many of the concerns raised by machine learning practitioners, and also introduced a number of new theoretical tools. Here we concentrate on applications of the extended model to the problem of obtaining upper bounds on sufficient training sample size. Further work will be required to obtain lower bounds on sample size needed, and to determine the computational complexity of finding decision rules with near minimal empirical risk. Some promising results along these lines are given in Kearns and Schapire (1990). However, even granting that such results can be obtained, the extended model still has a number of shortcomings in its present form. Some of these can be easily remedied, others may be more problematic.

First, we define the model only for a fixed decision rule space \mathcal{H} . The model should be extended to learning problems on a sequence of decision rule spaces $\{\mathcal{H}_n: n \geq 1\}$, where \mathcal{H}_n is a decision rule space on an n -attribute domain X_n (e.g., $[0, 1]^n$), and to families of decision rule spaces of different "complexities" on a fixed domain (Kearns *et al.*, 1987; Blumer *et al.*, 1989; Haussler *et al.*, 1991a), so that tradeoffs between decision rule complexity and empirical risk can be addressed. The former extension is easy, the latter more involved. One approach to the latter problem is via Vapnik's (1989) principle of structural risk minimization (see also Devroye, 1988). Other approaches include the *MDL* (see, e.g., Barron and Cover, 1990), regularization (see, e.g., Poggio and Girosi, 1989), and more general Bayesian methods (see, e.g., Berger, 1985).

Second, the constants in the upper bounds are still too large to give sample size estimates that are useful in practice. It may be difficult to improve them to the point where the results are directly usable in applied work. Thus even with matching asymptotic lower bounds, practitioners may still need to rely at least in part on empirically derived sample size bounds. It is possible that the Bayesian viewpoint may yield better tools for calculating sample complexities. Support for this belief is given in Clarke and Barron (1990), (manuscript), Haussler *et al.* (1991b), and Oppen and Haussler (1991a). However, necessary sample size estimates for decision rule spaces as general as those studied from the minimax perspective using uniform convergence have not yet been tackled from the Bayesian perspective.

Finally, many other issues would need to be considered in a complete treatment of the problem of overfitting, including distribution specific bounds on sample complexity (Theorem 2 is actually distribution specific, since the random covering numbers are distribution specific, yet we only apply it here in a distribution independent setting), decision rule spaces

with infinite pseudo and metric dimensions (these include various classes of “smooth” functions and their relatives; see Dudley, 1984, Chap. 7, and Quiroz, 1989), and non-i.i.d. sources of examples (see White, 1990a, and Nobel and Dembo, 1990). Despite these shortcomings, we feel that the theory we give here provides useful insights into the nature of the problem of overfitting in learning, and because of its generality will be a useful starting point for further research in this area.

9. APPENDIX

9.1. Metric Spaces, Covering Numbers, and Metric Dimension

A *pseudo metric* on a set S is a function ρ from $S \times S$ into \mathbb{R}^+ such that for all $x, y, z \in S$, $x = y \Rightarrow \rho(x, y) = 0$, $\rho(x, y) = \rho(y, x)$ (symmetry), and $\rho(x, z) \leq \rho(x, y) + \rho(y, z)$ (triangle inequality). If in addition $\rho(x, y) = 0 \Rightarrow x = y$, then ρ is a *metric*. (S, ρ) is a *(pseudo) metric space*. (S, ρ) is *complete* if every Cauchy sequence of points in S converges to a point in S ; (S, ρ) is *separable* if it contains a countable dense subset, i.e., a countable subset A such that for every $x \in X$ and $\varepsilon > 0$ there exists $a \in A$ with $\rho(x, a) < \varepsilon$. If $\rho(x, y) = 1 \Leftrightarrow x \neq y$ then ρ is called the *discrete metric*.

The *diameter* of a set $T \subseteq S$ is $\sup\{\rho(x, y) : x, y \in T\}$. If the diameter of T is finite then we say that T is *bounded*. For any $\varepsilon > 0$, an ε -*cover* for T is a finite set $N \subseteq S$ (not necessarily contained in T) such that for all $x \in T$ there is a $y \in N$ with $\rho(x, y) \leq \varepsilon$. If T has a (finite) ε -cover for all $\varepsilon > 0$ then T is *totally bounded*. (Note that this implies that (T, ρ) is separable and bounded.) In this case the function $\mathcal{N}(\varepsilon, T, \rho)$ denotes the size of the smallest ε -cover for T (w.r.t. the space S and the (pseudo) metric ρ). We refer to $\mathcal{N}(\varepsilon, T, \rho)$ as a *covering number*. A set $R \subseteq T$ is ε -*separated* if for all distinct $x, y \in R$, $\rho(x, y) > \varepsilon$. We denote by $\mathcal{M}(\varepsilon, T, \rho)$ the size of the largest ε -separated subset of T . We refer to $\mathcal{M}(\varepsilon, T, \rho)$ as a *packing number*. The third argument to \mathcal{N} and \mathcal{M} will be omitted when the metric ρ is clear from the context.

The following inequalities are easily verified (see, e.g., Kolmogorov and Tihomirov, 1961):

THEOREM 12. *If T is a totally bounded subset of the (pseudo) metric space (S, ρ) then for any $\varepsilon > 0$,*

$$\mathcal{M}(2\varepsilon, T, \rho) \leq \mathcal{N}(\varepsilon, T, \rho) \leq \mathcal{M}(\varepsilon, T, \rho).$$

Hence both these measures of boundedness, by covering number and by packing number, are equivalent to within a factor of 2 of ε . Following

Kolmogorov and Tihomirov (1961), we define the *upper metric dimension* of a (pseudo) metric space (S, ρ) by

$$\overline{\dim}(S) = \limsup_{\varepsilon \rightarrow 0} \frac{\log \mathcal{N}(\varepsilon, S, \rho)}{\log(1/\varepsilon)}.$$

The *lower metric dimension*, denoted by $\underline{\dim}$, of a (pseudo) metric space (S, ρ) , is defined similarly using \liminf . When $\overline{\dim}(S) = \underline{\dim}(S)$, then this quantity is denoted $\dim(S)$, and referred to simply as the *metric dimension* of (S, ρ) . This quantity has also been called the *fractal dimension* (Farmer, 1982) and the *capacity dimension* (Farmer *et al.*, 1983). A very lucid and intuitive treatment is given in Mandelbrot (1982).

9.2. Permissible Classes of Functions

In order to obtain the uniform convergence results given in Theorem 2, certain measurability assumptions have to be made concerning the class of functions \mathbf{F} when this class is uncountable. These we have indicated by saying that \mathbf{F} must be a *permissible* class (Pollard, 1984). Here we give a definition of permissible that is a special case of that given by Pollard. This definition will be suitable for our purposes; we refer the reader to Pollard (1984) and Dudley (1984) for a more general treatment. See Exercise 10, p. 39 of Pollard (1984) for an indication of the kind of problems that can come up with non-permissible classes.

Throughout the paper we have assumed that \mathbf{F} is a class of real-valued functions on a set Z , and that P is a measure defined on some σ -algebra \mathcal{A} of subsets of Z such that each function in \mathbf{F} is measurable. We need further conditions on \mathbf{F} when it is uncountable. Let us say that the class \mathbf{F} is *indexed* by the set T if

$$\mathbf{F} = \{f(\cdot, t): t \in T\},$$

where f is a real-valued function on $Z \times T$ and $f(\cdot, t)$ denotes the real-valued function on Z obtained from f by fixing the second parameter to t . We will say that the function f *indexes* \mathbf{F} by T .

We need some structure on T as well. If T is contained in a topological space \bar{T} then we say that T is a *Borel subspace* if T is a Borel set with respect to the topology on \bar{T} ; i.e., if T is in the smallest σ -algebra on \bar{T} containing the open sets. The σ -algebra $\mathcal{B}(T)$ of Borel sets on T is then the restriction to T of the σ -algebra \mathcal{B} of Borel sets on \bar{T} ; i.e.,

$$\mathcal{B}(T) = \{B \cap T: B \in \mathcal{B}\}.$$

Finally, if \mathcal{A} is any σ -algebra on Z and \mathcal{C} is any σ -algebra on T then $\mathcal{A} \times \mathcal{C}$ denotes the smallest σ algebra on $Z \times T$ that contains $\{A \times C: A \in \mathcal{A}, C \in \mathcal{C}\}$. We are now ready for our main definition.

DEFINITION 11. We say that the class \mathbf{F} is *permissible* if it can be indexed by a set T such that

1. T is a Borel subspace of a compact metric space \bar{T} and
2. the function $f: Z \times T \rightarrow \mathfrak{R}$ that indexes \mathbf{F} by T is measurable with respect to the σ -algebra $\mathcal{A} \times \mathcal{B}(T)$.

Most uncountable classes of functions that come up in practice can be indexed by a finite number of real parameters (i.e., with $T = \mathfrak{R}^n$ for some $n \geq 1$) in such a way that condition (2) is satisfied. Condition (1) is satisfied as well in this case, since we can take T to be the one-point compactification of T , obtained by adding a *point at infinity* to T (see, e.g., Simmons, 1963).

Results given in Pollard (1984, Appendix C) imply that the sets used in Lemmas 12 and 13 are measurable when \mathbf{F} is permissible. He also shows that the packing numbers $\mathcal{M}(\varepsilon, \mathbf{F}_{|\mathcal{Z}}, d_{L^1})$ are measurable functions of $\mathcal{Z} \in \mathcal{Z}^m$ for any $m \geq 1$. Since Theorem 12 relates these closely to the covering numbers $\mathcal{N}(\varepsilon, \mathbf{F}_{|\mathcal{Z}}, d_{L^1})$, this allows us to further formalize our usage of random covering numbers. A more formal treatment would either replace the covering numbers with the packing numbers in our upper bounds, or reword probabilistic bounds on the covering numbers to use outer measure arguments.

9.3. Measuring the Accuracy of Empirical Estimates with the d_v Metric

In this section we give two bounds on the probability of large deviation of empirical estimates from true means, as measured by the d_v metric. One is derived from Chebyshev's inequality and the other from Bernstein's inequality. The first bound is better for estimates obtained from small samples, the second for estimates obtained from larger samples.

LEMMA 9. Let Z_1, \dots, Z_n be i.i.d. random variables with range $0 \leq Z_i \leq M$ and $\mathbf{E}(Z_i) = \mu$, $1 \leq i \leq n$. Assume $v > 0$ and $0 < \alpha < 1$. Then

1. $\Pr\left(d_v\left(\frac{1}{n} \sum_{i=1}^n Z_i, \mu\right) > \alpha\right) \leq \frac{M^2}{4\alpha^2 v n (M + v)} < \frac{M}{4\alpha^2 v n}.$
2. $\Pr\left(d_v\left(\frac{1}{n} \sum_{i=1}^n Z_i, \mu\right) > \alpha\right)$

$$\leq 2e^{-18\alpha^2 v n / (3 + \alpha)^2 M} < 2e^{-(9/8)\alpha^2 v n / M} < 2e^{-\alpha^2 v n / M}.$$

Proof. Let $Y_i = Z_i - \mu$, $1 \leq i \leq n$. Then

$$\begin{aligned} \Pr \left(d_v \left(\frac{1}{n} \sum_{i=1}^n Z_i, \mu \right) > \alpha \right) \\ = \Pr \left(\frac{|\sum_{i=1}^n Z_i - \mu n|}{vn + \sum_{i=1}^n Z_i + \mu n} > \alpha \right) \\ \leq \Pr \left(\left| \sum_{i=1}^n Y_i \right| > \alpha n(v + \mu) \right), \end{aligned}$$

since $\sum_{i=1}^n Z_i \geq 0$.

To obtain the first bound, note that by Chebyshev's inequality,

$$\Pr \left(\left| \sum_{i=1}^n Y_i \right| > \alpha n(v + \mu) \right) \leq \sigma^2 / (\alpha n(v + \mu))^2,$$

where $\sigma^2 = n \text{Var}(Y_i)$. Since $0 \leq Z_i \leq M$, $\text{Var}(Z_i) = \text{Var}(Y_i) \leq \mu(M - \mu)$. Hence

$$\Pr \left(\left| \sum_{i=1}^n Y_i \right| > \alpha n(v + \mu) \right) \leq \frac{\mu(M - \mu)}{\alpha^2 n(v + \mu)^2}.$$

It is easily verified that the maximum value of this expression occurs at $\mu = (vM)/(2v + M)$, and that this gives an upper bound of

$$\frac{M^2}{4\alpha^2 v n(M + v)}.$$

To obtain the second bound, we apply Bernstein's inequality (see, e.g., Pollard, 1984, p. 192), which states that

$$\Pr \left(\left| \sum_{i=1}^n Y_i \right| > \eta \right) \leq 2e^{-\eta^2/2(n \text{Var}(Y_i) + (1/3)B\eta)}$$

for any zero mean i.i.d. random variables Y_1, \dots, Y_n bounded in absolute value by B . Substituting $\eta = \alpha n(v + \mu)$ and upper bounds $B \leq M$ and $\text{Var}(Y_i) \leq \mu M$, this gives a bound of

$$2e^{-\alpha^2 n^2 (v + \mu)^2 / 2(n\mu M + (1/3)M\alpha n(v + \mu))} = 2e^{-3\alpha^2 n(v + \mu)^2 / 2M(\alpha v + (3 + \alpha)\mu)}.$$

Since $(v + \mu)^2 / (\alpha v + (3 + \alpha)\mu)$ is minimized at $\mu = (3 - \alpha)/(3 + \alpha)v$, the latter expression is bounded by substituting this value of μ . This gives the first bound of part (2). The second bound follows from the fact that $\alpha < 1$. ■

9.4. Proof of Main Theorem on Uniform Convergence of Empirical Estimates

In this section we prove Theorem 3 from Section 3. The proof is given in a series of lemmas. We first extend the metric d_v to a pseudo metric on vectors in $(\mathfrak{R}^+)^{2m}$ for $m \geq 1$. We will do so in a somewhat unusual manner that will be useful in what follows. In fact, this extension can be defined for any metric, so we state it in its general form.

DEFINITION 12. For each integer $m \geq 1$, let Γ_{2m} denote the set of all permutations σ of $\{1, \dots, 2m\}$ such that for all i , $1 \leq i \leq m$, either $\sigma(i) = m + i$ and $\sigma(m + i) = i$, or $\sigma(i) = i$ and $\sigma(m + i) = m + i$. Thus the permutations in Γ_{2m} swap selected indices in the first half of the sequence $\{1, \dots, 2m\}$ with corresponding indices in the second half. For any $\vec{x} = (x_1, \dots, x_{2m}) \in (\mathfrak{R}^+)^{2m}$ and $\sigma \in \Gamma_{2m}$, let $\mu_1(\vec{x}, \sigma) = (1/m) \sum_{i=1}^m x_{\sigma(i)}$ and $\mu_2(\vec{x}, \sigma) = (1/m) \sum_{i=m+1}^{2m} x_{\sigma(i)}$. For any metric d on \mathfrak{R}^+ and $\vec{x}, \vec{y} \in (\mathfrak{R}^+)^{2m}$, let $\vec{d}(\vec{x}, \vec{y}) = \max\{d(\mu_1(\vec{x}, \sigma), \mu_1(\vec{y}, \sigma)) + d(\mu_2(\vec{x}, \sigma), \mu_2(\vec{y}, \sigma)) : \sigma \in \Gamma_{2m}\}$.

It is easily verified that \vec{d} is a pseudo-metric on $(\mathfrak{R}^+)^{2m}$. Symmetry is obvious, and the triangle inequality follows easily from the triangle inequality for d on \mathfrak{R}^+ :

$$\begin{aligned} \vec{d}(\vec{x}, \vec{y}) + \vec{d}(\vec{y}, \vec{z}) &= \max\{d(\mu_1(\vec{x}, \sigma), \mu_1(\vec{y}, \sigma)) + d(\mu_2(\vec{x}, \sigma), \mu_2(\vec{y}, \sigma)) : \sigma \in \Gamma_{2m}\} \\ &\quad + \max\{d(\mu_1(\vec{y}, \sigma), \mu_1(\vec{z}, \sigma)) + d(\mu_2(\vec{y}, \sigma), \mu_2(\vec{z}, \sigma)) : \sigma \in \Gamma_{2m}\} \\ &\geq \max\{d(\mu_1(\vec{x}, \sigma), \mu_1(\vec{y}, \sigma)) + d(\mu_2(\vec{x}, \sigma), \mu_2(\vec{y}, \sigma)) \\ &\quad + d(\mu_1(\vec{y}, \sigma), \mu_1(\vec{z}, \sigma)) + d(\mu_2(\vec{y}, \sigma), \mu_2(\vec{z}, \sigma)) : \sigma \in \Gamma_{2m}\} \\ &\geq \max\{d(\mu_1(\vec{x}, \sigma), \mu_1(\vec{z}, \sigma)) + d(\mu_2(\vec{x}, \sigma), \mu_2(\vec{z}, \sigma)) : \sigma \in \Gamma_{2m}\} \\ &= \vec{d}(\vec{x}, \vec{z}). \end{aligned}$$

We note the following additional property of this extension.

LEMMA 10. For all $\vec{x}, \vec{y} \in (\mathfrak{R}^+)^{2m}$ and $\sigma \in \Gamma_{2m}$, $d(\mu_1(\vec{x}, \sigma), \mu_2(\vec{x}, \sigma)) \leq d(\mu_1(\vec{y}, \sigma), \mu_2(\vec{y}, \sigma)) + \vec{d}(\vec{x}, \vec{y})$.

Proof. We have

$$\begin{aligned} d(\mu_1(\vec{x}, \sigma), \mu_2(\vec{x}, \sigma)) &\leq d(\mu_1(\vec{y}, \sigma), \mu_2(\vec{y}, \sigma)) + d(\mu_1(\vec{x}, \sigma), \mu_1(\vec{y}, \sigma)) \\ &\quad + d(\mu_2(\vec{x}, \sigma), \mu_2(\vec{y}, \sigma)) \end{aligned}$$

by the triangle inequality on d . The last two terms of this sum combined are at most $\vec{d}(\vec{x}, \vec{y})$ by definition. ■

We now restrict ourselves to the case that the metric d is the metric d_v for some $v > 0$. The following lemma will play a key role in establishing our basic exponential inequality.

LEMMA 11. *Let $\vec{x} = (x_1, \dots, x_{2m})$ be a sequence of reals such that $0 \leq x_i \leq M$, $1 \leq i \leq 2m$. Assume $v > 0$ and $0 < \alpha < 1$. Then if a permutation $\sigma \in \Gamma_{2m}$ is chosen uniformly at random,*

$$\Pr(d_v(\mu_1(\vec{x}, \sigma), \mu_2(\vec{x}, \sigma)) > \alpha) \leq 2e^{-2\alpha^2 vm/M}.$$

Proof. For each i , $1 \leq i \leq m$, let Y_i be an independent random variable such that $Y_i = x_i - x_{m+i}$ with probability $\frac{1}{2}$ and $Y_i = x_{m+i} - x_i$ with probability $\frac{1}{2}$. Note that for any $\sigma \in \Gamma_{2m}$,

$$\begin{aligned} d_v(\mu_1(\vec{x}, \sigma), \mu_2(\vec{x}, \sigma)) &= \frac{|(1/m) \sum_{i=1}^m x_{\sigma(i)} - (1/m) \sum_{i=m+1}^{2m} x_{\sigma(i)}|}{v + (1/m) \sum_{i=1}^{2m} x_{\sigma(i)}} \\ &= \frac{|\sum_{i=1}^m (x_{\sigma(i)} - x_{\sigma(m+i)})|}{vm + \sum_{i=1}^{2m} x_i}. \end{aligned}$$

Hence

$$\begin{aligned} &\Pr(d_v(\mu_1(\vec{x}, \sigma), \mu_2(\vec{x}, \sigma)) > \alpha) \\ &= \Pr\left(\left|\sum_{i=1}^m (x_{\sigma(i)} - x_{\sigma(m+i)})\right| > \alpha \left(vm + \sum_{i=1}^{2m} x_i\right)\right) \\ &= \Pr\left(\left|\sum_{i=1}^m Y_i\right| > \alpha \left(vm + \sum_{i=1}^{2m} x_i\right)\right), \end{aligned}$$

because each swap in a randomly chosen $\sigma \in \Gamma_{2m}$ is independent. Since $E(Y_i) = 0$ and $-|x_i - x_{m+i}| \leq Y_i \leq |x_i - x_{m+i}|$, we can apply Hoeffding's inequality (see, e.g., Pollard, 1984) to bound the latter probability by

$$2e^{-\alpha^2 (vm + \sum_{i=1}^{2m} x_i)^2 / 2 \sum_{i=1}^m (x_i - x_{m+i})^2}.$$

Let $\beta = \sum_{i=1}^{2m} x_i$. Since $0 \leq x_i \leq M$,

$$\sum_{i=1}^m (x_i - x_{m+i})^2 \leq \sum_{i=1}^m M |x_i - x_{m+i}| \leq \beta M.$$

Hence we have

$$2e^{-\alpha^2 (vm + \sum_{i=1}^{2m} x_i)^2 / 2 \sum_{i=1}^m (x_i - x_{m+i})^2} \leq 2e^{-\alpha^2 (vm + \beta)^2 / 2\beta M} = 2e^{-(\alpha^2 / 2M)(vm + \beta)^2 / \beta}.$$

The expression in parentheses is minimized, and therefore the whole expression is maximized, by setting $\beta = vm$, giving a value of $4vm$. Hence

$$\Pr(d_v(\mu_1(\vec{x}, \sigma), \mu_2(\vec{x}, \sigma)) > \alpha) \leq 2e^{-2\alpha^2 vm/M}. \quad \blacksquare$$

For our next lemma we need some notation to refer to the separate empirical estimates based on the first and second halves of an even length sample.

DEFINITION 13. For all $m \geq 1$ and $\vec{z} \in Z^{2m}$, we let $\hat{\mathbf{E}}'_z(f) = (1/m) \sum_{i=1}^m f(z_i)$ and $\hat{\mathbf{E}}''_z(f) = (1/m) \sum_{i=m+1}^{2m} f(z_i)$.

LEMMA 12. Let \mathbf{F} be a permissible set of functions on Z with $0 \leq f(z) \leq M$ for all $f \in \mathbf{F}$ and $z \in Z$. Assume $v > 0$, $0 < \alpha < 1$, and $m \geq 2M/(\alpha^2 v)$. Then

$$\begin{aligned} & \Pr\{\vec{z} \in Z^{2m}: \exists f \in \mathbf{F} \text{ with } d_v(\hat{\mathbf{E}}_z(f), \mathbf{E}(f)) > \alpha\} \\ & \leq 2\Pr\{\vec{z} \in Z^{2m}: \exists f \in \mathbf{F} \text{ with } d_v(\hat{\mathbf{E}}'_z(f), \hat{\mathbf{E}}''_z(f)) > \alpha/2\}. \end{aligned}$$

Proof. If Z and \mathbf{F} are uncountable, the assumption of permissibility guarantees that these probabilities are well-defined (see Section 9.2 and Pollard (1984)). From Chebyshev's inequality (see Lemma 9, part (1) in Section 9.3), for each individual $f \in \mathbf{F}$,

$$\Pr\{\vec{z} \in Z^{2m}: d_v(\hat{\mathbf{E}}_z(f), \mathbf{E}(f)) > \alpha/2\} \leq M/(\alpha^2 vm).$$

Since $m \geq 2M/(\alpha^2 v)$, this probability is at most $\frac{1}{2}$. Now consider any $f \in \mathbf{F}$ and sample $\vec{z}' \in Z^m$ such that $d_v(\hat{\mathbf{E}}_{\vec{z}'}(f), \mathbf{E}(f)) > \alpha$. If we draw an independent random sample $\vec{z}'' \in Z^m$, then with probability at least $\frac{1}{2}$, $d_v(\hat{\mathbf{E}}_{\vec{z}''}(f), \mathbf{E}(f)) \leq \alpha/2$. Whenever this happens we have $d_v(\hat{\mathbf{E}}_{\vec{z}'}(f), \hat{\mathbf{E}}_{\vec{z}''}(f)) > \alpha/2$ by the triangle inequality for d_v . Thus

$$\begin{aligned} & \Pr\{\vec{z} \in Z^{2m}: \exists f \in \mathbf{F} \text{ with } d_v(\hat{\mathbf{E}}'_z(f), \hat{\mathbf{E}}''_z(f)) > \alpha/2\} \\ & \geq \Pr\{\vec{z}' \vec{z}'' \in Z^{2m}: \exists f \in \mathbf{F} \text{ with } d_v(\hat{\mathbf{E}}_{\vec{z}'}(f), \mathbf{E}(f)) > \alpha \\ & \quad \text{and } d_v(\hat{\mathbf{E}}_{\vec{z}''}(f), \mathbf{E}(f)) \leq \alpha/2\} \\ & \geq \frac{1}{2} \Pr\{\vec{z}' \in Z^m: \exists f \in \mathbf{F} \text{ with } d_v(\hat{\mathbf{E}}_{\vec{z}'}(f), \mathbf{E}(f)) > \alpha\}. \end{aligned}$$

Again, when Z and \mathbf{F} are uncountable, permissibility guarantees that the implied use of Fubini's theorem in obtaining the above inequalities is justified. \blacksquare

We are now in a position to prove the following version of our theorem, using the extended metric \vec{d}_v in place of the L^1 metric to measure covering numbers.

LEMMA 13. Let \mathbf{F} be a permissible set of functions on Z with $0 \leq f(z) \leq M$ for all $f \in \mathbf{F}$ and $z \in Z$. Assume $v > 0$, $0 < \alpha < 1$, and $m \geq 1$. Let

$$p(\alpha, v, m) = \Pr\{\bar{z} \in Z^m: \exists f \in \mathbf{F} \text{ with } d_v(\hat{\mathbf{E}}_{\bar{z}}(f), \mathbf{E}(f)) > \alpha\}.$$

Then

$$p(\alpha, v, m) \leq 2\mathbf{E}(\min(2\mathcal{N}(\alpha/4, \mathbf{F}_{|\bar{z}}, \vec{d}_v) e^{-\alpha^2 vm/8M}, 1)),$$

and if in addition $\mathbf{F}_{|\bar{z}}$ is finite for all $\bar{z} \in Z^{2m}$ then

$$p(\alpha, v, m) \leq 2\mathbf{E}(\min(2|\mathbf{F}_{|\bar{z}}| e^{-\alpha^2 vm/2M}, 1)),$$

where the expectations are over \bar{z} drawn randomly from Z^{2m} .

Proof. First note that both bounds are trivial if $m < 2M/(\alpha^2 v)$, so we may assume $m \geq 2M/(\alpha^2 v)$. Hence by Lemma 12,

$$p(\alpha, v, m) \leq 2\Pr\{\bar{z} \in Z^{2m}: \exists f \in \mathbf{F} \text{ with } d_v(\hat{\mathbf{E}}'_{\bar{z}}(f), \hat{\mathbf{E}}''_{\bar{z}}(f)) > \alpha/2\}.$$

Thus it suffices to obtain bounds for the latter quantity.

We begin with the second bound, for the case when $\mathbf{F}_{|\bar{z}}$ is always finite. For any sample $\bar{z} = (z_1, \dots, z_{2m}) \in Z^{2m}$ and $\sigma \in \Gamma_{2m}$, let $\sigma(\bar{z}) = (z_{\sigma(1)}, \dots, z_{\sigma(2m)})$. For any fixed function $f \in \mathbf{F}$ and fixed $\bar{z} \in Z^{2m}$, if we select a permutation $\sigma \in \Gamma_{2m}$ uniformly at random,

$$\Pr(d_v(\hat{\mathbf{E}}'_{\sigma(\bar{z})}(f), \hat{\mathbf{E}}''_{\sigma(\bar{z})}(f)) > \alpha/2) \leq 2e^{-\alpha^2 vm/2M} \quad (7)$$

by Lemma 11. Hence for any fixed $\bar{z} \in Z^{2m}$, if we select a permutation $\sigma \in \Gamma_{2m}$ uniformly at random,

$$\Pr(\exists f \in \mathbf{F} \text{ with } d_v(\hat{\mathbf{E}}'_{\sigma(\bar{z})}(f), \hat{\mathbf{E}}''_{\sigma(\bar{z})}(f)) > \alpha/2) \leq \min(2|\mathbf{F}_{|\bar{z}}| e^{-\alpha^2 vm/2M}, 1). \quad (8)$$

Thus if we draw \bar{z} at random from Z^{2m} and independently select a permutation $\sigma \in \Gamma_{2m}$ uniformly at random,

$$\Pr(\exists f \in \mathbf{F} \text{ with } d_v(\hat{\mathbf{E}}'_{\sigma(\bar{z})}(f), \hat{\mathbf{E}}''_{\sigma(\bar{z})}(f)) > \alpha/2) \leq \mathbf{E}(\min(2|\mathbf{F}_{|\bar{z}}| e^{-\alpha^2 vm/2M}, 1)).$$

However, since each of the $2m$ observations in \bar{z} is independent, each of the samples $\sigma(\bar{z})$ for $\sigma \in \Gamma_{2m}$ is equally likely. Hence

$$\Pr(\exists f \in \mathbf{F} \text{ with } d_v(\hat{\mathbf{E}}'_{\sigma(\bar{z})}(f), \hat{\mathbf{E}}''_{\sigma(\bar{z})}(f)) > \alpha/2),$$

where both \bar{z} and σ are chosen at random, is the same as

$$\Pr(\exists f \in \mathbf{F} \text{ with } d_v(\hat{\mathbf{E}}'_{\bar{z}}(f), \hat{\mathbf{E}}''_{\bar{z}}(f)) > \alpha/2),$$

where only the sample \bar{z} is chosen at random. The second bound follows.

The proof of the first bound is similar, except for steps 7 and 8. From

Lemma 10, using the extension \vec{d}_v of the metric d_v to even-length sequences of reals given in Definition 12 above, if $f, f^* \in \mathbf{F}$ are such that

$$\vec{d}_v((f(z_1), \dots, f(z_{2m})), (f^*(z_1), \dots, f^*(z_{2m}))) \leq \alpha/4,$$

then for any $\sigma \in \Gamma_{2m}$ such that

$$d_v(\hat{\mathbf{E}}'_{\sigma(\vec{z})}(f), \hat{\mathbf{E}}''_{\sigma(\vec{z})}(f)) > \alpha/2,$$

we must have

$$d_v(\hat{\mathbf{E}}'_{\sigma(\vec{z})}(f^*), \hat{\mathbf{E}}''_{\sigma(\vec{z})}(f^*)) > \alpha/4.$$

Thus if N is an $\alpha/4$ -cover for $\mathbf{F}_{|\vec{z}}$ with respect to the metric \vec{d}_v , then whenever there exists $f \in \mathbf{F}$ with

$$d_v(\hat{\mathbf{E}}'_{\sigma(\vec{z})}(f), \hat{\mathbf{E}}''_{\sigma(\vec{z})}(f)) > \alpha/2,$$

there exists $f^* \in N$ with

$$d_v(\hat{\mathbf{E}}'_{\sigma(\vec{z})}(f^*), \hat{\mathbf{E}}''_{\sigma(\vec{z})}(f^*)) > \alpha/4.$$

For fixed f^* and random σ , the probability of the latter event is at most $2e^{-\alpha^2 vm/8M}$ by Lemma 11. Hence for any fixed $\vec{z} \in Z^{2m}$, if we select a permutation $\sigma \in \Gamma_{2m}$ uniformly at random,

$$\begin{aligned} & \Pr(\exists f \in \mathbf{F} \text{ with } d_v(\hat{\mathbf{E}}'_{\sigma(\vec{z})}(f), \hat{\mathbf{E}}''_{\sigma(\vec{z})}(f)) > \alpha/2) \\ & \leq \min(2\mathcal{N}(\alpha/4, \mathbf{F}_{|\vec{z}}, \vec{d}_v) e^{-\alpha^2 vm/8M}, 1). \end{aligned}$$

The remainder of the proof is as above. \blacksquare

We need only one more lemma to complete our proof, one that can be used to relate the \vec{d}_v covering numbers to the d_{L^1} covering numbers.

LEMMA 14. For any $m \geq 1$, $\vec{x}, \vec{y} \in (\mathbb{R}^+)^{2m}$, and $v > 0$, $\vec{d}_v(\vec{x}, \vec{y}) \leq (2/v) d_{L^1}(\vec{x}, \vec{y})$.

Proof. For any $\sigma \in \Gamma_{2m}$

$$\begin{aligned} & d_v(\mu_1(\vec{x}, \sigma), \mu_1(\vec{y}, \sigma)) + d_v(\mu_2(\vec{x}, \sigma), \mu_2(\vec{y}, \sigma)) \\ &= \frac{|\sum_{i=1}^m (x_{\sigma(i)} - y_{\sigma(i)})|}{vm + \sum_{i=1}^m (x_{\sigma(i)} + y_{\sigma(i)})} + \frac{|\sum_{i=m+1}^{2m} (x_{\sigma(i)} - y_{\sigma(i)})|}{vm + \sum_{i=m+1}^{2m} (x_{\sigma(i)} + y_{\sigma(i)})} \\ &\leq \frac{\sum_{i=1}^{2m} |x_{\sigma(i)} - y_{\sigma(i)}|}{vm} = \frac{2}{v} d_{L^1}(\vec{x}, \vec{y}). \end{aligned}$$

The result follows. \blacksquare

The theorem follows easily from the last two lemmas. \blacksquare

9.5. BOUNDS ON SAMPLE SIZE FOR LEARNING IN FEEDFORWARD NETS WITH SHARP THRESHOLDS

In this section we give the bounds for uniform convergence of empirical estimates in neural networks with sharp threshold functions claimed in Section 7.

THEOREM 13. *Let \mathcal{A} be a feedforward architecture as defined in Section 7 with $n \geq 1$ inputs, one output, $N \geq 2$ computation units, and a total of W weights and biases. Let $X = \mathbb{R}^n$, $Y = A = \{0, 1\}$, and l be the discrete loss function. Assume that the squashing function for each computation unit has the form $\text{sign} \circ \sigma$, where σ is non-decreasing or non-increasing. Different σ 's can be used for different units. Let \mathcal{H} be all functions from X into A representable on \mathcal{A} by varying the weights and biases. Let $v > 0$, $0 < \alpha < 1$ and $m \geq 1$. Suppose that \vec{z} is generated by m independent random draws from a probability measure P on $Z = X \times Y$. Then*

$$\Pr(\exists h \in \mathcal{H} : d_v(\hat{\mathbf{r}}_{h,l}(\vec{z}), \mathbf{r}_{h,l}(P)) > \alpha) \leq 4(2eNm/W)^W e^{-\alpha^2 vm/2}.$$

This probability is at most δ for a sample size m that is

$$O\left(\frac{1}{\alpha^2 v} \left(W \log \frac{N}{\alpha v} + \log \frac{1}{\delta}\right)\right).$$

Proof. Each computation unit in the network with k weights is associated with a class of $\{0, 1\}$ -valued functions of the form

$$f(\vec{x}) = \text{sign} \left(\sigma(\phi_1(\vec{x}), \dots, \phi_k(\vec{x})) + \theta + \sum_{j=1}^k w_j \phi_j(\vec{x}) \right),$$

where θ, w_1, \dots, w_k are adjustable real-valued parameters and $\phi_1, \dots, \phi_k, \mu$ and σ are fixed functions, the latter monotone. By Theorems 4 and 5 this class of functions has pseudo dimension at most $k+1$. Since the pseudo dimension is the same as the Vapnik-Chervonenkis dimension for classes of indicator functions, this implies that the class has Vapnik-Chervonenkis dimension at most $k+1$. Now let d be the sum of all the Vapnik-Chervonenkis dimensions of all the classes of functions associated with the computation units of the architecture \mathcal{A} . It follows that $d \leq W$, the total number of weights and biases in the network.

For each $h \in \mathcal{H}$ let l_h be the loss function associated with h for the discrete loss l , i.e., $l_h(x, y) = 1$ if $y \neq h(x)$, $l_h(x, y) = 0$ if $y = h(x)$. Let $\mathbf{F} = \mathbf{I}_{\mathcal{H}} = \{l_h : h \in \mathcal{H}\}$. Let $\vec{z} = ((x_1, y_1), \dots, (x_m, y_m))$ be any fixed sample and $\vec{x} = (x_1, \dots, x_m)$. It is easily verified that $|\mathbf{F}_{|\vec{z}}| = |\mathcal{H}_{|\vec{x}}|$. It is shown in Baum and Haussler (1989, Theorem 1), (and is also implied directly from

results in Cover (1968)) that for any class \mathcal{H} as above, $|\mathcal{H}_{|\vec{x}}| \leq (Nm/d)^d$ for all $\vec{x} = (x_1, \dots, x_m)$, where d and N are as above. This implies that

$$|\mathbf{F}_{|\vec{z}}| \leq (Nm/d)^d \leq (Nm/W)^W$$

for all samples \vec{z} of length m . Since each $l_h \in \mathbf{F}$ is a random variable that is bounded between 0 and 1, Theorem 3 (second part) shows that

$$\Pr(\exists h \in \mathbf{F}: d_v(\hat{\mathbf{E}}_{\vec{z}}(h), \mathbf{E}(h)) > \alpha) \leq 4(2eNm/W)^W e^{-\alpha^2 vm/2}.$$

This gives the first bound.

For the second bound, it can be shown that for sample size

$$m = \frac{4}{\alpha^2 v} \left(2W \ln \frac{16N}{\alpha^2 v} + \ln \frac{4}{\delta} \right)$$

we have

$$4(2eNm/W)^W e^{-\alpha^2 vm/2} \leq \delta.$$

To see this, first note that by rearranging, we get

$$\alpha^2 vm/2 \geq W \ln(2eNm/W) + \ln(4/\delta).$$

Thus it suffices to show that

$$\alpha^2 vm/4 \geq W \ln(2eNm/W) \quad \text{and} \quad \alpha^2 vm/4 \geq \ln(4/\delta).$$

The latter inequality is assured by the last term in the formula for m . For the former inequality, let us take m equal to the first term only, i.e.,

$$m = \frac{8W}{\alpha^2 v} \ln \frac{16N}{\alpha^2 v}.$$

Substituting this into the former inequality and simplifying, we get

$$2 \ln \frac{16N}{\alpha^2 v} \geq \ln \left(\frac{16eN}{\alpha^2 v} \ln \frac{16N}{\alpha^2 v} \right).$$

This further simplifies to

$$\frac{16N}{\alpha^2 v} \geq e \ln \frac{16N}{\alpha^2 v},$$

which holds, since $x \geq e \ln x$ for any x . Finally, since this inequality holds for the given m , it is easy to see that it will also hold for larger m . ■

ACKNOWLEDGMENTS

I thank Dana Angluin, David Pollard, and Phil Long for their careful criticisms of an earlier draft of this paper, and their numerous suggestions for improvements. I also thank Naoki Abe, Anselm Blumer, Richard Dudley, and Michael Kearns for helpful comments on earlier drafts. I also thank Ron Rivest, David Rumelhart, Andrzej Ehrenfeucht, and Nick Littlestone for stimulating discussions on these topics.

RECEIVED October 5, 1989; FINAL MANUSCRIPT RECEIVED March 18, 1992

REFERENCES

- ABE, N., AND WARMUTH, M. (1990), On the computational complexity of approximating distributions by probabilistic automata, in "Proceedings of the 3rd Workshop on Computational Learning Theory," pp. 52–66, Morgan Kaufmann, San Mateo.
- ALEXANDER, K. (1985), Rates of growth for weighted empirical processes, in "Proceedings of Berkeley Conference in Honor of Jerzy Neyman and Jack Kiefer," Vol. 2, pp. 475–493.
- ALEXANDER, K. (1987), Rates of growth and sample moduli for weighted empirical processes indexed by sets, *Probab. Theory Related Fields* **75**, 379–423.
- ANGLUIN, D. (1988), Queries and concept learning, *Mach. Learning* **2**, 319–342.
- ANGLUIN, D., AND LAIRD, P. (1988), Learning from noisy examples, *Mach. Learning* **2** (4), 343–370.
- ANGLUIN, D., AND VALIANT, L. G. (1979), Fast probabilistic algorithms for Hamiltonian circuits and matchings, *J. Comput. System Sci.* **18**, 155–193.
- ANTHONY, M., AND SHAW-TAYLOR, J. (1990), "A result of Vapnik with Applications," Technical Report CSD-TR-628, University of London, Surrey.
- ASSOUAD, P. (1983), Densité et dimension, *Ann. Inst. Fourier* **33** (3), 233–282.
- BARRON, A. (1989), Statistical properties of artificial neural networks, in "28th Conference on Decision and Control," pp. 280–285.
- BARRON, A., AND COVER, T. (1990), Minimum complexity density estimation, *IEEE Trans. Inform. Theory*, to appear.
- BARTO, A. G., AND ANANDAN, P. (1985), Pattern recognizing stochastic learning automata, *IEEE Trans. Systems Man Cybernet.* **15**, 360–374.
- BAUM, E., AND HAUSSLER, D. (1989), What size net gives valid generalization? *Neural Comput.* **1** (1), 151–160.
- BENEDEK, G. M., AND ITAI, A. (1988), Learnability by fixed distributions, in "Proceedings, 1988 Workshop on Comp. Learning Theory," pp. 80–90, Morgan Kaufmann, San Mateo, CA.
- BERGER, J. (1985), "Statistical Decision Theory and Bayesian Analysis," Springer-Verlag, New York.
- BILLINGSLEY, P. (1986), "Probability and Measure," Wiley, New York.
- BLUMER, A., EHRENFUCHT, A., HAUSSLER, D., AND WARMUTH, M. K. (1989), Learnability and the Vapnik–Chervonenkis dimension, *J. Assoc. Comput. Mach.* **36** (4), 929–965.
- BREIMAN, L., FRIEDMAN, J. H., OLSEN, R. A., AND STONE, C. J. (1984), "Classification and Regression Trees," Wadsworth.
- BUNTINE, W. L. (1990), "A Theory of Learning Classification Rules," Ph.D. Thesis, University of Technology, Sydney.
- BUNTINE, W., AND WEIGEND, A. (1991), Bayesian back propagation, unpublished manuscript.
- CLARKE, B., AND BARRON, A. (manuscript), Entropy, risk and the Bayesian central limit theorem.

- CLARKE, B., AND BARRON, A. (1990), Information-theoretic asymptotics of Bayes methods, *IEEE Trans. Inform. Theory* **36** (3), 453–471.
- COVER, T. M. (1968), Capacity problems for linear machines, in “Pattern Recognition” (L. Kanal, Ed.), pp. 283–289. Thompson.
- DENKER, J., SCHWARTZ, D., WITTNER, B., SOLLA, S., HOWARD, R., JACKEL, L., AND HOPFIELD, J. (1987), Automatic learning, rule extraction and generalization, *Complex Systems* **1**, 877–922.
- DEVROYE, L. (1988), Automatic pattern recognition: A study of the probability of error, *IEEE Trans. Pattern Anal. Mach. Intelligence* **10** (4), 530–543.
- DUDA, R. O., AND HART, P. E. (1973), “Pattern Classification and Scene Analysis,” Wiley, New York.
- DUDLEY, R. M. (1978), Central limit theorems for empirical measures, *Ann. Probab.* **6** (6), 899–929.
- DUDLEY, R. M. (1984), A course on empirical processes, in “Lecture Notes in Mathematics,” Vol. 1097, pp. 2–142, Springer-Verlag, Berlin/New York.
- DUDLEY, R. M. (1987), Universal Donsker classes and metric entropy, *Ann. Probab.* **15** (4), 1306–1326.
- DURBIN, R., AND RUMELHART, D. E. (1989), Product units: A computationally powerful and biologically plausible extension to backpropagation networks, *Neural Comput.* **1** (1), 133–142.
- EDELSBRUNNER, H. (1987), “Algorithms in Combinatorial Geometry,” Springer-Verlag, Berlin/New York.
- EHRENFEUCHT, A., HAUSSLER, D., KEARNS, M., AND VALIANT, L. (1989), A general lower bound on the number of examples needed for learning, *Inform. Comput.* **82**, 247–261.
- FARMER, J. D. (1982), Information dimension and the probabilistic structure of chaos, *Z. Naturforsch. A* **37**, 1304–1325.
- FARMER, J. D., OTT, E., AND YORKE, J. A. (1983), The dimension of chaotic attractors, *Physica D* **7**, 153–180.
- FERGUSON, T. (1967), “Mathematical Statistics: A Decision Theoretic Approach,” Academic Press, New York.
- GULLAPALLI, V. (1990), A stochastic reinforcement algorithm for learning real-valued functions, *Neural Networks* **3** (6), 671–692.
- GYORGY, G., AND TISHBY, N. (1990), Statistical theory of learning a rule, in “Neural Networks and Spin Glasses” (K. Thuemann and R. Koeberle, Eds.), World Scientific.
- HAUSSLER, D. (1988), Quantifying inductive bias: AI learning algorithms and Valiant’s learning framework, *Artificial Intelligence* **36**, 177–221.
- HAUSSLER, D. (1989), Learning conjunctive concepts in structural domains, *Mach. Learning*, **4**, 7–40.
- HAUSSLER, D. (1990), Decision theoretic generalizations of the PAC learning model, in “Proceedings, First Workshop on Algorithmic Learning Theory, Tokyo, Japan,” pp. 21–41.
- HAUSSLER, D., KEARNS, M., LITTLESTONE, N., AND WARMUTH, M. K. (1991a), Equivalence of models for polynomial learnability, *Inform. Comput.* **95**, 129–161.
- HAUSSLER, D., KEARNS, M., AND SCHAPIRE, R. (1991b), Bounds on the sample complexity of Bayesian learning using information theory and the VC dimension, in “Proceedings of the Fourth Workshop on Computational Learning Theory,” pp. 61–74.
- HAUSSLER, D., LITTLESTONE, N., AND WARMUTH, M. (1990), “Predicting $\{0, 1\}$ -Functions on Randomly Drawn Points,” Technical Report UCSC-CRL-90-54, Computer Research Laboratory, University of California, Santa Cruz; *Inform. Comput.*, to appear.
- HAUSSLER, D., AND WELZL, E. (1987), Epsilon nets and simplex range queries, *Discrete Comput. Geom.* **2**, 127–151.
- KEARNS, M., LI, M., PITT, L., AND VALIANT, L. (1987), On the learnability of Boolean

- formulae, in "Proceedings of the Nineteenth Annual ACM Symposium on Theory of Computing," pp. 285–295, New York.
- KEARNS, M., AND SCHAPIRE, R. (1990), Efficient distribution-free learning of probabilistic concepts, in "31st Annual IEEE Symposium on Foundations of Computer Science," pp. 382–391.
- KIEFER, J. (1987), "Introduction to Statistical Inference," Springer-Verlag, Berlin/New York.
- KOLMOGOROV, A. N., AND TIHOMIROV, V. M. (1961), ϵ -entropy and ϵ -capacity of sets in functional spaces, *Amer. Math. Soc. Transl. Ser. 2* 17, 277–364.
- KULKARNI, S. (1989), "On Metric Entropy, Vapnik–Chervonenkis Dimension, and Learnability for a Class of Distributions," Technical Report LIDS-P-1910, Center for Intelligent Control Systems, MIT.
- KULLBACK, S. (1959), "Information Theory and Statistics," Wiley, New York.
- LECUN, Y., DENKER, J., AND SOLLA, S. (1990), Optimal brain damage, in "Advances in Neural Information Processing Systems" (D. Touretsky, Ed.), Vol. 2, p. 589, Morgan Kaufmann, San Mateo, CA.
- LINDLEY, D. V. (1990), The present position in Bayesian statistics, *Statist. Sci.* 5 (1), 44–89.
- LINEAL, N., MANSOUR, Y., AND RIVEST, R. (1988), Results on learnability and the Vapnik–Chervonenkis dimension, in "Proceedings of the 1988 Workshop on Computational Learning Theory," pp. 56–68, Morgan Kaufmann, San Mateo, CA.
- LITTLESTONE, N. (1988), Learning quickly when irrelevant attributes abound: A new linear-threshold algorithm, *Mach. Learning* 2, 285–318.
- MACKEY, D. (1992), "Bayesian Methods for Adaptive Models," Ph.D. Thesis, California Institute of Technology.
- MANDELBROT, B. B. (1982), "The Fractal Geometry of Nature," Freeman, San Francisco.
- MCCULLAGH, P., AND NELDER, J. A. (1989), "Generalized Linear Models," Chapman & Hall, London.
- MOODY, J., AND DARKEN, C. (1989), Fast learning in networks of locally-tuned processing units, *Neural Comput.* 1 (2), 281–294.
- NARENDRA, K. S., AND THATHACHAR, M. A. L. (1989), "Learning Automata—An Introduction," Prentice-Hall, Englewood Cliffs, NJ.
- NATARAJAN, B. K. (1988), Learning over classes of distributions, in "Proceedings of the 1988 Workshop on Computational Learning Theory," pp. 408–409, Morgan Kaufmann, San Mateo, CA.
- NATARAJAN, B. K. (1989a), "Probably-Approximate Learning over Classes of Distributions." Technical Report HPL-SAL-89-29, Hewlett Packard Labs, Palo Alto, CA.
- NATARAJAN, B. K. (1989b), "Some Results on Learning," Technical Report CMU-RI-TR-89-6, Carnegie Mellon.
- NATARAJAN, B. K., AND TADEPALLI, P. (1988), Two new frameworks for learning, in "Proceedings of the 5th International Conference on Machine Learning," pp. 402–415, Morgan Kaufmann, San Mateo, CA.
- NOBEL, A., AND DEMBO, A. (1990), "On Uniform Convergence for Dependent Processes." Technical Report 74, Dept. of Statistics, Stanford University, Stanford, CA.
- NOLAN, D., AND POLLARD, D. (1987), U-processes: Rates of convergence, *Ann. Statist.* 15 (2), 780–799.
- NOWLAN, S. (1990), Maximum Likelihood Competitive Learning, in "Advances in Neural Information Processing Systems" (D. Touretsky, Ed.), Vol. 2, pp. 574–582, Morgan Kaufmann, San Mateo, CA.
- NOWLAN, S., AND HINTON, G. (1991), "Soft Weight-Sharing," Technical Report, Department of Computer Science, University of Toronto.
- OPPER, M., AND HAUSSLER, D. (1991a), Calculation of the learning curve of Bayes optimal classification algorithm for learning a perceptron with noise, in "Computational Learning

- Theory: Proceedings of the Fourth Annual Workshop," pp. 75–87, Morgan Kaufmann, San Mateo, CA.
- OPPER, M., AND HAUSSLER, D. (1991b), Generalization performance of Bayes optimal classification algorithm for learning a perceptron, *Phys. Rev. Lett.* **66** (20), 2677–2680.
- POGGIO, T., AND GIROSI, F. (1989), "A Theory of Networks for Approximation and Learning," Technical Report A.I. Memo No. 1140, Massachusetts Institute of Technology, Cambridge, MA.
- POLLARD, D. (1984), "Convergence of Stochastic Processes," Springer-Verlag, Berlin/New York.
- POLLARD, D. (1986), Rates of uniform almost-sure convergence for empirical processes indexed by unbounded classes of functions, manuscript.
- POLLARD, D. (1990), "Empirical Processes: Theory and Applications," NSF–CBMS Regional Conference Series in Probability and Statistics, Vol. 2, Inst. of Math. Stat. and Am. Stat. Assoc.
- QUIROZ, A. J. (1989), Metric entropy and learnability, unpublished manuscript, Universidad Simón Bolívar, Caracas, Venezuela.
- RENYI, A. (1970), "Probability Theory," North-Holland, Amsterdam.
- RISSANEN, J. (1986), Stochastic complexity and modeling, *Ann. Statist.* **14** (3), 1080–1100.
- RUMELHART, D. E., AND MCCLELLAND, J. L. (1986), "Parallel Distributed Processing: Explorations in the Microstructure of Cognition, Volume 1: Foundations," MIT Press, Cambridge, MA.
- SAUER, N. (1972), On the density of families of sets, *J. Combin. Theory Ser. A* **13**, 145–147.
- SHACKELFORD, G., AND VOLPER, D. (1988), Learning k -DNF with noise in the attributes, in "Proceedings of the 1988 Workshop on Computational Learning Theory," pp. 97–103, Morgan Kaufmann, San Mateo, CA.
- SIMMONS, G. F. (1963), "Introduction to Topology and Modern Analysis," McGraw-Hill, New York.
- SLOAN, R. (1988), Types of noise in data for concept learning, in "Proceedings, 1988 Workshop on Comp. Learning Theory," pp. 91–96, Morgan Kaufmann, San Mateo, CA.
- SOMPOLINSKY, H., TISHBY, N., AND SEUNG, H. S. (1990), Learning from examples in large neural networks, *Phys. Rev. Lett.* **65**, 1683–1686.
- TALAGRAND, M. (1991), Sharper bounds for empirical processes, unpublished manuscript.
- TISHBY, N., LEVIN, E., AND SOLLA, S. (1989), Consistent inference of probabilities in layered networks: Predictions and generalizations, in "IJCNN International Joint Conference on Neural Networks," Vol. II, pp. 403–409. IEEE, New York.
- TOURETSKY, D. (1989), "Advances in Neural Information Processing Systems," Vol. 1, Morgan Kaufmann, San Mateo, CA.
- TOURETSKY, D. (1990), "Advances in Neural Information Processing Systems," Vol. 2, Morgan Kaufmann, San Mateo, CA.
- VALIANT, L. G. (1984), A theory of the learnable, *Comm. ACM* **27** (11), 1134–1142.
- VAPNIK, V. N. (1982), "Estimation of Dependences Based on Empirical Data," Springer-Verlag, New York.
- VAPNIK, V. N. (1989), Inductive principles of the search for empirical dependences (methods based on weak convergence of probability measures), in "Proceedings of the 2nd Workshop on Computational Learning Theory," Morgan Kaufmann, San Mateo, CA.
- VAPNIK, V. N., AND CHERVONENKIS, A. YA., On the uniform convergence of relative frequencies of events to their probabilities, *Theory Probab. Appl.* **16** (2), 264–280.
- WEIGEND, A., HUBERMAN, B., AND RUMELHART, D. (1990), Predicting the future: A connectionist approach, *Internat. J. Neural Systems* **1**, 193–209.
- WEISS, S., AND KULIKOWSKI, C. (1991), "Computer Systems That Learn," Morgan Kaufmann, San Mateo, CA.

- WELZL, E. (1988), Partition trees for triangle counting and other range search problems, in "4th ACM Symposium on Computational Geometry," pp. 23–33, Urbana, IL.
- WENOCUR, R. S., AND DUDLEY, R. M. (1981), Some special Vapnik–Chervonenkis classes, *Discrete Math.* **33**, 313–318.
- WHITE, H. (1990a), Connectionist nonparametric regression: Multilayer feedforward networks can learn arbitrary mappings, *Neural Networks* **3**, 535–549.
- WHITE, H. (1990b), Learning in artificial neural networks: A statistical perspective, *Neural Comput.* **1** (4), 425–464.
- YAMANISHI, K. (1990), A learning criterion for stochastic rules, in "Proceedings of the 3rd Workshop on Computational Learning Theory," pp. 67–81, Morgan Kaufmann.